



Antimicrobial Resistance and Biofilm Encoding Genes Amongst the *Staphylococcus aureus* Bacteria Isolated From Meat and Meat Products



CrossMark

Kobra Abbasi, Elahe Tajbakhsh* and Hassan Momtaz

Department of Microbiology, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran.

ROLE of meat and meat products as reservoirs of biofilm formation and antimicrobial-resistant *Staphylococcus aureus* is not clearly observed. The present research was conducted to assess the incidence, antimicrobial resistance and distribution of *icaABCD* biofilm formation genes amongst the *S. aureus* bacteria recovered from raw meat and meat products. One-hundred and sixty raw meat and meat product samples were collected and *S. aureus* bacteria were isolated using microbial culture. Antimicrobial resistance of *S. aureus* bacteria was assessed by disk diffusion. PCR was used to assess the distribution of *icaABCD* genes. Twenty-six out of 160 (16.25%) examined samples were contaminated with *S. aureus*. Raw beef meat (40%) and raw chicken burger (12.50%) samples had the highest *S. aureus* contamination rate. Bacteria harbored the highest resistance toward penicillin (69.23%), ampicillin (61.53%), tetracycline (57.69%) and gentamicin (57.69%) antimicrobials. *IcaD*, *icaA*, *icaB* and *icaC* genes were detected in 65.38%, 57.69%, 50%, and 42.30% *S. aureus* bacteria, respectively. Roles of raw beef, chicken, beef burger and chicken burger samples as reservoirs of antimicrobial resistant-and biofilm formation-*S. aureus* bacteria have been demonstrated in the current research. Nevertheless, supplementary investigations should perform to understand other roles of *icaABCD* biofilm formation genes in *S. aureus* originated from meat and meat products.

Keywords, *Staphylococcus aureus*; Incidence; Antibiotic resistance; Biofilm formation; Meat.

Introduction

Staphylococcus aureus (*S. aureus*) a pathogenic bacterium accountable for emerging foodborne diseases globally. Food related outcomes of the staphylococcal diseases characterized by nausea, weakness, abdominal cramps, vomiting, and toxic shock syndrome (TST) [1,2]. Contaminated food products, predominantly those originated from animals (especially meat and meat products), are known as *S. aureus* reservoirs [3-5].

Staphylococcal foodborne diseases are mostly treated with antimicrobial-based therapeutic options. Nevertheless, newly-launched researches announced the high incidence of resistance

of *S. aureus* bacteria originated from animal resources toward diverse classes of antimicrobial agents, particularly cephalosporins, penicillin, macrolides, aminoglycosides, tetracyclines, and fluoroquinolones [6,7].

From a molecular view, biofilm formation is one of the most critical ways of *S. aureus* bacteria to become virulence and resist toward diverse antimicrobial agents [8]. Polysaccharide intracellular adhesion (PIA) factor, which is encoded by the *ica* genes (*icaA*, *icaB*, *icaC* and *icaD*). These genes are involved in biofilm activities and caused intracellular adhesion and protected *S. aureus* bacteria from immune response and antimicrobial agents [8].

Rendering the boost importance of biofilm formation *S. aureus* bacteria as a potential food-borne pathogen, The current survey was performed to assess the antimicrobial resistance and frequency of biofilm formation genes of the *S. aureus* bacteria recovered from meat and meat products.

Materials and Methods

Sampling

In the current cross sectional descriptive survey, a total of 160 raw meat and meat product samples including beef (n= 40) and chicken (n= 40) meat and meat burger (n= 40) and chicken burger (n= 40) samples were randomly collected from supermarkets of Shahrekord city, Iran. Meat samples (100 g) were collected from the femur muscle. Meat product samples (100 g) were collected from valid brands. Samples were transferred to research center using small refrigerator.

Isolation and identification of S. aureus

Twenty-five grams of each sample were merged with 225 mL of buffered peptone water (Merck, Germany) and homogenized by the Stomacher device (Interscience, Saint-Nom, France). At that point, 5 mL of the achieved sample was inoculated on 50 mL Trypticase Soy Broth (TSB, Merck, Germany) medium (with 10% NaCl and 1% sodium pyruvate). Media were incubated at 35 °C for 18 h. At that point, a loopful of an achieved culture was inoculated on the Baird-Parker agar (BPA, Merck, Germany) (contained egg yolk tellurite emulsion). Media were incubated at 37 °C for about 24 h. *S. aureus* colonies were determined as a black shiny color with 2 to 5-mm clear marginal zones. *S. aureus* identification was accompanied using standard tests including Gram-staining, catalase, oxidase, mannitol fermentation, DNase activity, and sugar fermentation tests [6].

Antimicrobial resistance determination

Guidelines of the Clinical Laboratory Standard Institute (CLSI) [9] were applied to determine the antimicrobial resistance pattern of isolated *S. aureus* bacteria. *S. aureus* bacteria were cultured on the Muller Hinton Agar (MHA, Merck, Germany) media and antimicrobial disks (penicillin (10 µg/disk) (P10), tetracycline (30 µg/disk) (T30), methicillin (5 µg/disk) (Met5), ampicillin (10 µg/disk) (AM10), oxacillin (1 µg/disk) (Ox1), vancomycin (5 µg/disk) (V5), gentamicin (10 µg/disk) (G10), rifampin (5 µg/

disk) (Rif5) and erythromycin (15 µg/disk) (Er15) (Himedia, India)) were placed on plate media. Media contained *S. aureus* bacteria (0.5 McFarland concentration) and disks were incubated at 37 °C for 24 h. After incubation, diameters of the growth inhibition zone surround the bacteria were measured and compared with those of CLSI. Diameters of the growth inhibition zones of bacteria were compared with those presented in CLSI and then resistance or sensitive strains were determined. *S. aureus* (ATCC 29213) was cultured as control.

Determination of biofilm formation encoding-genes

S. aureus isolates were sub-cultured on TSB and incubated for 48 h at 37 °C. Genomic DNA was extracted from the colonies using the DNA extraction kit (Thermo Fisher Scientific, Germany) rendering the guideline. Quality and quantity of extracted DNA were assessed by nanodrop device (NanoDrop, Thermo Scientific, USA) and electrophoresis (2% agarose gel). Table 1 described the list of primers and PCR circumstances applied for detection of biofilm formation genes [10, 11].

DNA thermo-cycler (Eppendorf, Germany) was applied. Fifteen microliters of the amplified samples were subjected to gel electrophoresis (2% gel contained 0.1% ethidium bromide (0.4 µg/ml)). Results of electrophoresis were checked by the UVdoc (London, UK). *S. aureus* (ATCC 29213) and PCR -grade water (Thermo Fisher Scientific, Germany) were applied as positive and negative controls, respectively[6].

Statistic

Data obtained from the research were transferred to Excel software and analyzed by SPSS 21.0 software (Chicago, USA) using Chi-square and Fisher's exact tests. *P* value <0.05 was determined as significant level.

Results

Table 2 described the incidence of *S. aureus* amongst examined meat and meat product samples. Twenty-six out of 160 (16.25%) examined samples were contaminated with *S. aureus*. Beef (40%) had the higher contamination rate than chicken (5%) (*P*<0.05). Chicken burger (12.50%) had the higher contamination rate than beef burger (7.50%). However, it was not statistically significant (*P*>0.05).

TABLE 1. Primers and PCR circumstances applied for detection of biofilm formation genes [10, 11].

Target genes	Primer sequence (5'-3')	PCR product (bp)	PCR programs	PCR volume (50µL)
<i>icaA</i>	F, ACACTTGCTGGCGCAGTCAA R, TCTGGAACCAACATCCAACA	188	1 cycle, 94 ^{0C} ----- 5 min. 30 cycles, 94 ^{0C} ----- 60 s 55 ^{0C} ----- 60 s 72 ^{0C} ----- 60 s 1 cycle, 72 ^{0C} ----- 10 min	
<i>icaB</i>	F, AGAATCGTGAAGTATAGAAAATT R, TCTAATCTTTTCATGGAATCCGT	900	1 cycle, 94 ^{0C} ----- 5 min. 30 cycles, 94 ^{0C} ----- 60 s 52 ^{0C} ----- 30 s 72 ^{0C} ----- 90 s 1 cycle, 72 ^{0C} ----- 10 min	5 µL PCR buffer 10X 1.5 mM MgCl ₂ 200 µM dNTP (Fermentas) 0.5 µM of each primers F & R
<i>icaC</i>	F, ATGGGACGGATTCCATGAAAAAGA R, TAATAAGCATTAATGTTCAATT	1100	1 cycle, 94 ^{0C} ----- 5 min. 30 cycles, 94 ^{0C} ----- 60 s 55 ^{0C} ----- 30 s 72 ^{0C} ----- 30 s 1 cycle, 72 ^{0C} ----- 10 min	1.25 U Taq DNA polymerase (Fermentas) 2.5 µL DNA template
<i>icaD</i>	F, ATGGTCAAGCCCAGACAGAG R, AGTATTTTCAATGTTTAAAGCAA	198	1 cycle, 94 ^{0C} ----- 5 min. 25 cycles, 94 ^{0C} ----- 30 s 55 ^{0C} ----- 30 s 72 ^{0C} ----- 60 s 1 cycle, 72 ^{0C} ----- 10 min	

DNA thermo-cycler (Eppendorf, Germany) was applied. Fifteen microliters of the amplified samples were subjected to gel electrophoresis (2% gel contained 0.1% ethidium bromide (0.4 µg/ml)). Results of electrophoresis were checked by the UVdoc (London, UK). *S. aureus* (ATCC 29213) and PCR -grade water (Thermo Fisher Scientific, Germany) were applied as positive and negative controls, respectively[6].

Table 3 described the antimicrobial resistance pattern of *S. aureus* isolates. *S. aureus* isolates exhibited the uppermost incidence of resistance toward penicillin (69.23%), ampicillin (61.53%), tetracycline (57.69%) and gentamicin (57.69%). Statistically significant variance was obtained between types of samples and incidence of antimicrobial resistance ($P < 0.05$).

Table 4 describes the incidence of biofilm formation genes amongst the *S. aureus* bacteria

recovered from meat and meat products. Total distribution of *icaD*, *icaA*, *icaB* and *icaC* amongst the *S. aureus* isolates were 65.38%, 57.69%, 50%, and 42.30%, respectively. *IcaD* had the uppermost incidence amongst biofilm formation genes, while *icaC* had the lowermost. *S. aureus* bacteria recovered from both beef and chicken burger samples had the higher distribution of biofilm formation genes than those of beef and chicken ($P < 0.05$).

TABLE 2. Incidence of *S. aureus* amongst examined meat and meat product samples.

Samples	N. collected	N (%). Positive for <i>S. aureus</i>
Beef	40	16 (40)
Chicken	40	2 (5)
Beef burger	40	3 (7.50)
Chicken burger	40	5 (12.50)
Total	160	26 (16.25)

TABLE 3. Antimicrobial resistance pattern of *S. aureus* isolates recovered from meat and meat products.

Samples (N. <i>S. aureus</i>)	N. (%) <i>S. aureus</i> harbored resistance to each antibiotic								
	P10	T30	Met5	AM10	Ox1	V5	G10	Rif5	Er15
Beef (16)	12 (75)	9 (56.25)	9 (56.25)	10 (62.50)	8 (50)	4 (25)	9 (56.25)	7 (43.75)	7 (43.75)
Chicken (2)	1 (50)	1 (50)	1 (50)	1 (50)	1 (50)	-	1 (50)	1 (50)	1 (50)
Beef burger (3)	2 (66.66)	2 (66.66)	2 (66.66)	2 (66.66)	1 (33.33)	1 (33.33)	2 (66.66)	1 (33.33)	1 (33.33)
Chicken burger (5)	3 (60)	3 (60)	2 (40)	3 (60)	2 (40)	1 (20)	3 (60)	2 (40)	2 (40)
Total (26)	18 (69.23)	15 (57.69)	14 (53.84)	16 (61.53)	12 (46.15)	6 (23.07)	15 (57.69)	11 (42.30)	11 (42.30)

TABLE 4. Incidence of biofilm formation genes amongst the *S. aureus* bacteria recovered from meat and meat products.

Samples (N. <i>S. aureus</i>)	N. (%) <i>S. aureus</i> harbored each gene			
	<i>icaA</i>	<i>icaB</i>	<i>icaC</i>	<i>icaD</i>
Beef (16)	8 (50)	7 (43.75)	5 (31.25)	10 (62.50)
Chicken (2)	1 (50)	1 (50)	-	1 (50)
Beef burger (3)	2 (66.66)	2 (66.66)	2 (66.66)	2 (66.66)
Chicken burger (5)	4 (80)	3 (60)	4 (80)	4 (80)
Total (26)	15 (57.69)	13 (50)	11 (42.30)	17 (65.38)

Discussion

The importance of biofilm formation *S. aureus* as an emerging resistant bacteria in foodstuffs, particularly those with animal origins, is not documented well. Nevertheless, growing consideration has been focused on understanding staphylococcal biofilms as imperative factors for the occurrence of antimicrobial resistance [12].

An existing research was completed to assess the antimicrobial resistance pattern and distribution of biofilm formation genes amongst the *S. aureus* bacteria recovered from meat and meat products. Findings revealed that 16.25% of examined samples were contaminated with *S. aureus*. Incidence of *S. aureus* bacteria amongst raw meat samples examined in surveys conducted on Denmark [13], Egypt [14], Turkey [15], Germany [16] and Brazil [17] were 52%, 40.80%, 30%, 71.50% and 21.72%, respectively. Incidence of *S. aureus* in meat products in Iran [18], China [19], India [20], USA [21], Iraq [22], and Nigeria [23] were 9.86%, 35%, 21.81%, 27.90%, 54.20%, and 9.70%, respectively. Total incidence of *S. aureus* amongst the beef, camel meat, chicken, mutton, beef burger, beef sausage, ground beef, chicken burger, ground chicken, and chicken sausage samples collected from Libya were 20%, 23.80%, 40%, 0%, 25%, 33%, 36.40%, 50%, 20%, and 10%, respectively [24]. Brazilian survey [25] revealed that the incidence of *S. aureus* amongst the beef burger, beef sandwich, chicken burger and chicken sandwich samples were 64%, 20%, 72% and 8%, respectively. Variances in types of samples, method of experiment, geographical and seasonal circumstances may cause such difference in the incidence of *S. aureus* in diverse researches. Primary presence of *S. aureus* in meat samples through slaughtering and transmission of *S. aureus* from workers and staff of the slaughterhouses, veterinarians and butchers are two imperative routes of meat and meat products contamination.

S. aureus isolates harbored the considerable incidence of resistance toward penicillin, ampicillin, tetracycline and gentamicin which may partly be owing to the high prescription of these antimicrobial agents. Additionally, irregular and boost use of these antimicrobial, particularly in aviculture and veterinary caused significant occurrence of antimicrobial resistance. Excessive use of disinfectants, self-medication with antimicrobials and administration of single-dose therapeutics may be other probable reasons.

Similar to our findings, boost incidence rate of resistance of *S. aureus* bacteria toward penicillin, ampicillin, tetracycline and gentamicin was reported from China [19], India [26], South Africa [27], Iran [28], Pakistan [29], and United Kingdom [30]. Similar to our survey, Fowoyo and Ogunbanwo [31] described that the considerable incidence of resistance of *S. aureus* bacteria recovered from foodstuffs toward erythromycin (15.70%), oxacillin (35.70%), ampicillin (86.70%), gentamicin (11.40%), and trimethoprim-sulfamethoxazole (74.90%). Diverse views of veterinary physicians about type of prescribed antimicrobials, viability of rules about limitation of uses of antimicrobials, and prices of antimicrobial agents may affect the incidence of antimicrobial resistance in different regions.

High distribution of *icaABCD* biofilm formation genes increased the importance of isolated *S. aureus* strains. The *icaABCD* genes encrypt a Polysaccharide Intercellular Adhesion (PIA), which causes severe protection of *S. aureus* bacteria toward contrary environmental circumstances such as presence of antimicrobials and antiseptic agents and also immune responses. Furthermore, the *ica* biofilm formation gene acts as an imperative factors in nosocomial infections [32] and foodborne diseases [33, 34]. In a similar survey conducted by Ferreira, Tette, Mendonça, Soares and Carvalho [35], *icaA* and *icaD* genes were detected in 97% of the *S. aureus* bacteria recovered from poultry products. It seems that the *S. aureus* carrying *icaABCD* biofilm formation genes may be more important because of its ability to cause antimicrobial-resistant diseases. However, its role as an important foodborne agent has been overlooked. Diverse kinds of food-borne bacteria are responsible for food poisoning due to the consumption of raw or undercooked meat and meat products [36-42].

Conclusion

Boost incidence of *S. aureus* resists toward diverse antimicrobials and carrying *icaABCD* biofilm formation genes was reported in the current survey. High incidence of resistance toward routinely used antimicrobials and high distribution of *icaABCD* biofilm formation genes may reflect emerging public health issues rendering the consumption of contaminated beef, chicken, beef burger and chicken burger in raw or undercooked circumstances. However, further researches are mandatory to obtain other

epidemiological importance of biofilm formation *S. aureus* in meat and meat products.

Acknowledgements

Many thanks from staffs of the Food Hygiene Research Unit, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran.

Conflict of interest : The authors declared that no conflict of interest.

Funds statement : Self-funding

References

1. Abdolmaleki, Z., Mashak, Z. and Dehkordi, F.S., Phenotypic and genotypic characterization of antibiotic resistance in the methicillin-resistant *Staphylococcus aureus* strains isolated from hospital cockroaches. *Antimicrob. Resist. Infect. Control.*, **8**(1),1-14 (2019).
2. Abdolmaleki, Z., Mashak, Z. and Safarpour Dehkordi, F., Molecular and Virulence Characteristics of Methicillin-Resistant *Staphylococcus aureus* Bacteria Recovered From Hospital Cockroaches. *Jundishapur. J. Microbiol.*, **12**(12),e98564 (2019).
3. Rahi, A., Kazemeini, H., Jafariaskari, S., Seif, A., Hosseini, S. and Dehkordi, F.S., Genotypic and Phenotypic-Based Assessment of Antibiotic Resistance and Profile of Staphylococcal Cassette Chromosome mec in the Methicillin-Resistant *Staphylococcus aureus* Recovered from Raw Milk. *Infect. Drug. Resist.*, **13**,273–283 (2020).
4. Hasanpour Dehkordi, A., Khaji, L., Sakhaei Shahreza, M., Mashak, Z., Safarpour Dehkordi, F., Safae, Y., Hosseinzadeh, A., Alavi, I., Ghasemi, E. and Rabiei-Faradonbeh, M., One-year prevalence of antimicrobial susceptibility pattern of methicillin-resistant *Staphylococcus aureus* recovered from raw meat. *Trop. Biomed.*, **34**(2),396-404 (2017).
5. Madahi, H., Rostami, F., Rahimi, E. and Dehkordi, F.S., Prevalence of enterotoxigenic *Staphylococcus aureus* isolated from chicken nugget in Iran. *Jundishapur. J. Microbiol.*, **7**(8), e10237 (2014).
6. Safarpour Dehkordi, F., Gandomi, H., Basti, A.A., Misaghi, A. and Rahimi, E., Phenotypic and genotypic characterization of antibiotic resistance of methicillin-resistant *Staphylococcus aureus* isolated from hospital food. *Antimicrob. Resist. Infect. Control.*, **6**(1),Article number,104, pages 1-13(2017).
7. Momtaz, H., Dehkordi, F.S., Rahimi, E., Asgarifar, A. and Momeni, M., Virulence genes and antimicrobial resistance profiles of *Staphylococcus aureus* isolated from chicken meat in Isfahan province, Iran. *J. App. Poultry. Res.*, **22**(4),913-921 (2013).
8. Nourbakhsh, F. and Namvar, A.E., Detection of genes involved in biofilm formation in *Staphylococcus aureus* isolates. *GMS. Hyg. Infect. Control.*, **11**,1-5 (2016).
9. CLSI, Performance Standards for Antimicrobial Susceptibility Testing: Twenty-Fifth Informational Supplement, *CLSI document M100-S25*. (2015).
10. Deighton, M.A., Capstick, J., Domalewski, E. and Van Nguyen, T., Methods for studying biofilms produced by *Staphylococcus epidermidis*. *Methods in enzymology, Elsevier*; (2001).
11. Arciola, C.R., Campoccia, D., Gamberini, S., Cervellati, M., Donati, E. and Montanaro, L., Detection of slime production by means of an optimised Congo red agar plate test based on a colourimetric scale in *Staphylococcus epidermidis* clinical isolates genotyped for *ica* locus. *Biomaterial.*, **23**(21),4233-4239 (2002).
12. Rodríguez-Lázaro, D., Alonso-Calleja, C., Oniciuc, E.A., Capita, R., Gallego, D., González-Machado, C., Wagner, M., Barbu, V., Eiros-Bouza, J.M. and Nicolau, A.I., Characterization of biofilms formed by foodborne methicillin-resistant *Staphylococcus aureus*. *Front. Microbiol.*, **9**, Article number,3004, pages 1-12(2018).
13. Tang, Y., Larsen, J., Kjeldgaard, J., Andersen, P.S., Skov, R. and Ingmer, H., Methicillin-resistant and-susceptible *Staphylococcus aureus* from retail meat in Denmark. *Int. J. Food. Microbiol.*, **249**,72-76 (2017).
14. Karmi, M., Prevalence of methicillin-resistant *Staphylococcus aureus* in poultry meat in Qena, Egypt. *Vet. World.*, **6**(10),711-715 (2013).
15. Gundogan, N., Citak, S., Yucel, N. and Devren, A., A note on the incidence and antibiotic resistance of *Staphylococcus aureus* isolated from meat and chicken samples. *Meat. Sci.*, **69**(4), 807-810 (2005).
16. Richter, A., Sting, R., Popp, C., Rau, J., Tenhagen, B.-A., Guerra, B., Hafez, H. and Fetsch, A., Prevalence of types of methicillin-resistant *Staphylococcus aureus* in turkey flocks and personnel attending the animals. *Epidemiol. Infect.*, **140**(12),2223-2232 (2012).

17. Costa, W.L.R., Ferreira, J.d.S., Carvalho, J.S., Cerqueira, E.S., Oliveira, L.C. and Almeida, R.C.d.C., Methicillin-resistant *Staphylococcus aureus* in raw meats and prepared foods in public hospitals in Salvador, Bahia, Brazil. *J. Food. Sci.*, **80**(1),M147-M150 (2015).
18. Pourbabaee, M., Hadadi, M.R., Hooshyar, H., Pourbabaee, P. and Nazari-Alam, A., Prevalence of *Staphylococcus aureus* in raw hamburgers from Kashan in 2017. *Int. Arch. Health. Sci.*, **7**(1),47-50 (2020).
19. Wu, S., Huang, J., Wu, Q., Zhang, J., Zhang, F., Yang, X., Wu, H., Zeng, H., Chen, M. and Ding, Y., *Staphylococcus aureus* isolated from retail meat and meat products in China, incidence, antibiotic resistance and genetic diversity. *Front. Microbiol.*, **9**, Article number, 2767, pages 1-14 (2018) doi, [10.3389/fmicb.2018.02767](https://doi.org/10.3389/fmicb.2018.02767).
20. Zehra, A., Gulzar, M., Singh, R., Kaur, S. and Gill, J., Prevalence, multidrug resistance and molecular typing of methicillin-resistant *Staphylococcus aureus* (MRSA) in retail meat from Punjab, India. *J. Global. Antimicrob. Resist.*, **16**,152-158 (2019).
21. Ge, B., Mukherjee, S., Hsu, C.-H., Davis, J.A., Tran, T.T.T., Yang, Q., Abbott, J.W., Ayers, S.L., Young, S.R. and Creney, E.T., MRSA and multidrug-resistant *Staphylococcus aureus* in US retail meats, 2010–2011. *Food. Microbiol.*, **62**,289-297 (2017).
22. Kanaan, M.H., Antibacterial effect of ozonated water against methicillin-resistant *Staphylococcus aureus* contaminating chicken meat in Wasit Province, Iraq. *Vet. World.*, **11**(10), 1445-1453 (2018).
23. Ndahi, M., Kwaga, J., Bello, M., Kabir, J., Umoh, V., Yakubu, S. and Nok, A., 'Prevalence and antimicrobial susceptibility of *Listeria monocytogenes* and methicillin-resistant *Staphylococcus aureus* strains from raw meat and meat products in Zaria, Nigeria. *Letter. App. Microbiol.*, **58**(3),262-269 (2014).
24. Naas, H.T., Edarhoby, R.A., Garbaj, A.M., Azwai, S.M., Abolghait, S.K., Gammoudi, F.T., Moawad, A.A., Barbieri, I. and Eldaghayes, I.M., Occurrence, characterization, and antibiogram of *Staphylococcus aureus* in meat, meat products, and some seafood from Libyan retail markets. *Vet. World.*, **12**(6),925-931 (2019).
25. Contreras, C.P.Á., da Silva, L.N.N., Ferreira, D.C.G., dos Santos Ferreira, J. and de Castro Almeida, R.C., Prevalence of methicillin-resistant *Staphylococcus aureus* in raw hamburgers and ready-to-eat sandwiches commercialized in supermarkets and fast food outlets in Brazil. *Food. Nutr. Sci.*, **6**(14),1324-1331 (2015).
26. Savariraj, W.R., Ravindran, N.B., Kannan, P., Paramasivam, R., Senthikumar, T., Kumarasamy, P. and Rao, V.A., Prevalence, antimicrobial susceptibility and virulence genes of *Staphylococcus aureus* isolated from pork meat in retail outlets in India. *J. Food. Safe.*, **39**(1),e12589 (2019).
27. Fri, J., Njom, H.A., Ateba, C.N. and Ndip, R.N., Antibiotic Resistance and Virulence Gene Characteristics of Methicillin-Resistant *Staphylococcus aureus* (MRSA) Isolated from Healthy Edible Marine Fish. *Int. J. Microbiol.*, **2020**,1-9 (2020).
28. Rahimi, F. and Karimi, S., Characteristics of methicillin resistant *Staphylococcus aureus* strains isolated from poultry in Iran. *Arch. Clin. Infect. Dis.*, **10**(4),e30885 (2015).
29. Akbar, A. and Anal, A.K., Prevalence and antibiogram study of *Salmonella* and *Staphylococcus aureus* in poultry meat. *Asian. Pacific. J. Trop. Biomed.*, **3**(2),163-168 (2013).
30. Anjum, M.F., Marco-Jimenez, F., Duncan, D., Marín, C., Smith, R.P. and Evans, S., Livestock-associated Methicillin-Resistant *Staphylococcus aureus* from animals and animal products in the UK. *Front. Microbiol.*, **10**, Article number, 2136, pages 1-7(2019) doi.org/10.3389/fmicb.2019.02136.
31. Fowoyo, P. and Ogunbanwo, S., Antimicrobial resistance in coagulase-negative staphylococci from Nigerian traditional fermented foods. *Ann. Clin. Microbiol. Antimicrob.*, **16**(4),1-7 (2017).
32. Piechota, M., Kot, B., Frankowska-Maciejewska, A., Gruzewska, A. and Woźniak-Kosek, A., Biofilm formation by methicillin-resistant and methicillin-sensitive *Staphylococcus aureus* strains from hospitalized patients in Poland. *BioMed. Res. Int.*, **2018**,1-7 (2018).

33. Miao, J., Lin, S., Soteyome, T., Peters, B.M., Li, Y., Chen, H., Su, J., Li, L., Li, B. and Xu, Z., Biofilm formation of *Staphylococcus aureus* under food heat processing conditions, first report on CML production within biofilm. *Sci. Report.*, **9**(1),1-9 (2019).
34. Avila-Novoa, M.-G., Iñiguez-Moreno, M., Solís- Velázquez, O.-A., González-Gómez, J.-P., Guerrero-Medina, P.-J. and Gutiérrez-Lomelí, M., Biofilm formation by *Staphylococcus aureus* isolated from food contact surfaces in the dairy industry of Jalisco, Mexico. *J. Food. Quality.*, **2018**,1-8 (2018).
35. Ferreira, A.A., Tette, P.A.S., Mendonça, R.C.S., Soares, A.d.S. and Carvalho, M.M.D., Detection of exopolysaccharide production and biofilm-related genes in *Staphylococcus* spp. isolated from a poultry processing plant. *Food. Sci. Techno.*, **34**(4),710-716 (2014).
36. Momtaz, H., Davood Rahimian, M. and Safarpour Dehkordi, F., Identification and characterization of *Yersinia enterocolitica* isolated from raw chicken meat based on molecular and biological techniques. *J. App. Poultry. Res.*, **22**(1),137-145 (2013).
37. Rahimi, E., Yazdanpour, S. and Dehkordi, F., Detection of *Toxoplasma gondii* antibodies in various poultry meat samples using enzyme linked immuno sorbent assay and its confirmation by polymerase chain reaction. *J. Pure. Appl. Microbio.*, **8**(1),421-427 (2014).
38. Dehkordi, F., Parsaei, P., Saberian, S., Moshkelani, S., Hajshafiei, P., Hosseini, S., Babaei, M. and Ghorbani, M., Prevalence study of *Theileria annulata* by comparison of four diagnostic techniques in southwest Iran. *Bulgar. J. Vet. Med.*, **15**,123-130 (2012).
39. Ghorbani, F., Gheisari, E. and Dehkordi, F.S., Genotyping of *vacA* alleles of *Helicobacter pylori* strains recovered from some Iranian food items. *Tropical Journal of Pharmaceutical Research*, **15**(8),1631-1636 (2016).
40. Ranjbar, R., Masoudimanesh, M., Dehkordi, F.S., Jonaidi-Jafari, N. and Rahimi, E., Shiga (Vero)-toxin producing *Escherichia coli* isolated from the hospital foods; virulence factors, o-serogroups and antimicrobial resistance properties. *Antimicrob. Resist. Infect. Control.*, **6**(1),1-11(2017).
41. Hemmatinezhad, B., Khamesipour, F., Mohammadi, M., Safarpour Dehkordi, F. and Mashak, Z., Microbiological Investigation of O-Serogroups, Virulence Factors and Antimicrobial Resistance Properties of Shiga Toxin-Producing *Escherichia Coli* Isolated from Ostrich, Turkey and Quail Meats. *J. Food. Safe*, **35**(4),491-500 (2015).
42. Momtaz, H., Dehkordi, F.S., Rahimi, E., Ezadi, H. and Arab, R., Incidence of Shiga toxin-producing *Escherichia coli* serogroups in ruminant's meat. *Meat. Sci.*, **95**(2),381-388 (2013).
43. Ranjbar, R., Seif, and A. Dehkordi, F.S., Incidence of Antibiotic Resistance and Distribution of Virulence Factors in the Shiga Toxigenic *Escherichia coli* Recovered from Hospital Food. *Jundishapur.J. Microb.*, **12**(5),1-8 (2019).