



## In Vivo Study of Physiological, and Histological Effects of Different Doses of Mercury Oxide on Liver and Kidney in Male Wistar Rats

Abdulsatar A. Haji<sup>1\*</sup>, Alan I. Yousif<sup>1</sup>, Chinar M. Mohammed<sup>1</sup>, Thamer M. Bashir<sup>1</sup> and Ihsan H. Mohammed<sup>2</sup>

<sup>1</sup> Biology Department, Faculty of Science, the University of Zakho, Zakho, Kurdistan Region, Iraq.

<sup>2</sup> Medical Physiology and Pharmacology Department, College of Medicine, the University of Duhok, Duhok, Kurdistan Region, Iraq.



CrossMark

### Abstract

MERCURY accumulation affects the gastrointestinal, and renal systems. In this study, we aimed to study the physiological, and histological effects of mercury oxide on the liver and kidney in male Wistar rats. During 22 days, we divided 25 rats into 5 groups. The control group is placed first, followed by vinegar, low, medium, and high dose mercury groups. The control group was given only water. The vinegar-only group was given only vinegar. Mercury oxide-treated (HgO) group was given HgO 0.375 mg/kg/day. Mercury oxide treated group given HgO 1.5 mg/kg/day. Mercury oxide-treated (HgO) group was given HgO 4.5 mg/kg/day. We studied the levels of ALP, LDH, AST, ALT, albumin, creatinine, and urea. Histopathology of the liver and kidney were also studied. The result of this study was hepatic sinusoid dilation, renal tubule degeneration, and glomerulus shrinkage. This study showed non-significant differences among groups in terms of renal glomerulus diameter. The results showed that HgO at dose (1.5 mg/kg/day) had significantly higher levels of LDH, ALT, and AST enzymes when compared to the control group. While at the highest dose of mercury oxide (4.5 mg/kg/day), LDH, ALT, and AST enzyme levels decreased when compared to the lower doses. Our results showed a non-significant increase in urea level. Consequently, our investigation demonstrated that exposure to mercury oxide after therapy may result in toxicity to the kidneys and liver.

**Keywords:** Mercury oxide, Toxicity, Histopathology, Liver, Kidney.

### Introduction

Mercury oxide (HgO) is an inorganic compound that consists mainly of one atom of oxygen (O) and one atom of mercury (Hg). In nature, there are two main forms of mercury oxide, red and yellow. The most important component of the red form of mercury oxide is mercury. Mercury oxide is a very toxic heavy metal [1, 2] and it is known to pose a critical environmental hazard [3, 4]. This compound has countless industrial applications. It is used in medical measurement instruments, pesticides, dyes and fertilizers [5], cosmetics, glass modifiers, antiseptic compounds, [6], and batteries production [7, 8, 9].

In the year 2013, an agreement was signed among 147 countries around the world to regulate mercury global releases. Despite this, globally, minor emissions and pollution of mercury take place [10]. Mercury has been shown to cause a genotoxicity effect, due to its ability to bind sulfhydryl groups [11]. In many countries, as a result of its serious toxicity, its usage has been restricted.

Mercury is much more widely distributed than other heavy metals because of its high mobility [12, 13]. However, its effects on human health and the environment have long been documented. Because of its extensive usage, it had a noteworthy

\*Corresponding authors: Abdulsatar A. Haji, E-mail: [Abdulsatar.haji@staff.uoz.edu.krd](mailto:Abdulsatar.haji@staff.uoz.edu.krd) Tel.: 07502153874

(Received 16 July 2024, accepted 02 September 2024)

DOI: 10.21608/EJVS.2024.303733.2259

©National Information and Documentation Center (NIDOC)

influence on human health [11]. Many ways mercury-contaminated humans and one of the most important ways is via inhaling the vapor of mercury from gold mining, forest fires, and volcanic eruptions [14, 15, 16, 17]. Another way of mercury contamination is via consumption of contaminated fish [1, 2], which can cause life-threatening health problems, involving nephrotoxic [18], pneumotoxic [19, 20, 21], hepatotoxicity [19, 21, 22], cardiovascular and digestive systems toxicity [23, 25, 26, 27, 28]. Although, to date, there is doubt related to the relationship between cancer development and mercury exposure [11]. Mercury oxide like mercury, is extremely toxic and considered a worldwide transported pollutant. While there are many studies on mercury toxicity in human health, to our knowledge, there has been limited research on the *in vivo* effect of mercury oxide on the liver and kidney. So, the main aim of this study is to measure the physiological, and histopathology effects of different doses of mercury oxide on the liver and kidney of male albino rats.

## **Material and Methods**

### *Animals*

For this study, male healthy Wister rats (*Rattus norvegicus*) weighing 200 g – 300 g were used. Animals were kept under standard laboratory conditions of 12/12 hours dark/light cycle and room temperature (25 °C) with free access to water and food in clean cages in the Animal House of the University of Zakho.

### *Preparation of Mercury Oxide Doses*

Mercury oxide was dissolved in vinegar. For preparation doses of mercury oxide, we dissolved, 0.1125 mg of mercury oxide in 0.5ml of vinegar to obtain the first dose. For the second dose, we dissolved 0.45 mg of mercury oxide in 0.5 ml vinegar. For the third dose, 1.35 mg of mercury oxide was dissolved in 0.5 ml vinegar.

### *Experimental Design*

Each group with five rats a total of 25 rats were divided into 5 groups. Group 1 which is the untreated control group, was gavaged pure water. Group 2 which is the vinegar-treated group, was gavaged pure vinegar. Group 3, was gavaged the first dose of mercury oxide (0.375 mg/kg/day). Group 4, was gavaged the second dose of mercury oxide (1.5 mg/kg/day). Group 5, was gavaged the third dose of mercury oxide (4.5 mg/kg/day) for 22 days.

### *Physiological and Histopathological Studies*

After obtaining approval from the University of Zakho Animal Research Ethics Committees. Chloroform was used to anesthetize all rats. All rats were dissected to take out directly 5 ml of fresh non-coagulated blood from the hearts by using 5 ml medical disposable syringes. Immediately, collected blood samples were divided into two types of blood tubes, EDTA and plane tube. Then, all collected blood samples were sent to the laboratory. At that moment, the liver and right kidneys of all rats were removed, cleaned with distilled water, and weighed. Subsequently, they fixed in 10% of neutral buffered formalin, ascending grade of ethyl alcohol was used to dehydrate kidneys, cleared in xylene, and fixed in paraffin wax. They sectioned at a size of 5  $\mu$ m and hematoxylin and eosin stain were used to stain them. Sections were examined at 400x magnification using a compound microscope and a Dino-Eye microscopic camera was used to measure the diameter of 10 randomly chosen glomeruli in the cortex of examined kidneys of each rat. In the same way, the pathology of the liver was examined.

### *Statistical analysis*

For statistical analysis, GraphPad Prism Version 9 software was used. One-way analysis of variance (ANOVA) and the Dunnett test were used to compare variances among groups. A *P-value*  $\leq 0.05$  is considered significant among groups. Results are presented as means  $\pm$  standard error [24].

## **Results**

### *The Body Weight*

Our results reveal that body weight rates of all treated groups of rats significantly decreased after 22 days of exposure to mercury oxide compared to the control group (Fig.1). I.e., when the rats were gavaged with mercury oxide at a dose of 1.5 mg/kg/day, the body weight rates decreased to  $238.5 \pm 9.8$  g, compared to  $287.7 \pm 85.2$  g for the control group.

### *The Enzymes Activities of the Kidneys*

A minor increase occurred in urea levels in the kidneys of rats given large doses of mercury oxide compared to the control group (Table 1). In contrast, the levels of urea were decreased when low dosages of mercury oxide 0.375 mg/kg/day were given. Nevertheless, when compared to the higher dose and control, the changes are not significant. Compared to the control group, the

creatinine levels remained fairly higher at various mercury oxide dosages. I.e., creatinine levels of the control group  $0.3 \pm 0.01$  U/l were raised to  $0.4 \pm 0.051$  U/l at dose 0.375 mg/kg/day and reached  $0.35 \pm 0.03$  U/l at dose 4.5 mg/kg/day. That is, the changes are not significant.

#### *The Enzymes Activities of the Livers*

Our data suggest that there are significant differences in AST, ALT, ALP, and LDH, (*P-value* 0.0092, 0.0476, 0.0034, and 0.0194, respectively) levels between the control and mercury oxide-treated groups (Table 2). In contrast, our results show that there is a non-significant difference in GGT, albumin, and protein levels (*P-value* 0.121, 0.056, and 0.218 respectively) between the control and treated groups. When we compared the lower dose (0.375 mg/kg/day) to the higher dose (4.5 mg/kg/day) treated mercury oxide groups, the maximum dose showed a decrease in the AST, ALT, ALP, and LDH levels. In the mercury oxide treated (1.5 mg/kg/day) group, the AST, ALT, ALP, and LDH enzyme levels were significantly increased compared to the higher dose (4.5 mg/kg/day). Our data shows significant differences when we compared albumin levels of the control group, to the dose 1.5 mg/kg/day.

#### *Histopathology of Kidneys*

When the control and vinegar-treated kidneys of rats were examined (Figures 2 A and B), the renal glomeruli were found to have normal Bowman's space, normal capsules, and normal mesangial cells, proximal and distal convoluted tubules. Mercury oxide administration caused various observed effects on the renal morphology including, glomeruli size reduction and fragmentation into bifurcated components as well as renal tubules and glomeruli degeneration and blood extravasation in the medulla (Figures 2 C and D, and Figure 3). However, we did not observe any significant differences in the weights of kidneys and the diameter of glomeruli among the control and the experimental groups (Figure 4 and Figure 5).

#### *Histopathology of Liver*

The liver of the control and vinegar-treated groups showed a normal histological structure of the hepatocytic plate, hepatic sinusoid, and central vein (Figures 6 A and B). Although the hepatocyte structures are normal, mercury oxide administration in group 3 and group 4 resulted in noticeable alterations in the liver structure. These

changes included the dilation of a hepatic sinusoid (Figures 6 C and D). We observed a significant difference in liver weights between the control and treated groups (Figure 7).

#### **Discussion**

Globally heavy metals are accumulated in the environment. Especially, highly toxic mercury. Its accumulation in the environment increased gradually because it was not banned from use in industries (29). Studies showed that humans around the world are exposed to mercury (30). In a study on animals, mercury exposure caused loss of appetite and severe weight loss. This harmful effect may inhibit several vital metabolic processes within the body and eventually may lead to delays in development and growth (31).

There is a strong link between environmental metal exposure and chronic kidney diseases. As kidneys are very susceptible to the toxic effects of metals (32). Mercury exposure may cause adverse effects, such as nephrotoxicity, hepatotoxicity, neurotoxicity, immunotoxicity, teratogenicity, and cardiovascular and endocrine toxicity (33-34).

To date, the exact mechanism by which mercury oxide may cause nephrotoxicity is not clear. However, studies indicate that the reactive oxygen species (ROS) act as an important kidney disease mediator. Metabolism of mercury oxide in cells produced ROS such as hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), singlet oxygen (1O<sub>2</sub>), hydroxyl radical (.OH), superoxide anion (O<sub>2</sub><sup>-</sup>), and peroxy radicals (HOO. -). ROS toxicity is caused by antioxidant defense system disruption which can lead to damage to cellular DNA, proteins, and lipids (35).

According to studies heavy metals have serious side effects on mammalian organs (36). Our data indicated that there are significant increases in serum AST, ALT, ALP, and LDH levels. This data reinforces the data in the study done by Zaki *et al.* in 2011, which found significant increases in serum AST, ALT, and ALP levels in mercury oxide-treated catfish. The cytotoxic effects of mercury oxide may be attributed to glutathione, metallothionein, and protease activity alteration. Moreover, it is known that mercury oxide can produce ROS that can cause an increase in lipid peroxidation, which sequentially leads to a reduction of cell membrane integrity and eventually, cell death. Also, cell death may be caused by failure of DNA repair systems (37). Mercury oxide can disturb cellular growth,

proliferation, and differentiation processes and may cause some enzymes to be inactivated, and others like caspase to be activated and may cause alterations in the ultrastructure of hepatocytes (33). Mercury oxide-induced apoptosis in the liver may be caused by epigenetic mechanisms (34). Additionally, mercury oxide-induced behavior changes may occur in the brain due to neurotransmitter modulation including serotonin and dopamine (38). Currently, we discussed some mechanisms of mercury oxide-induced toxicities, but many are still far from being clearly understood.

### Conclusion

From the results we obtained from this study, we conclude that mercury oxide should be considered a major relevant risk factor for kidney and liver diseases. One potential limitation

of our work as we only studied two organs, the kidney and liver of rats, and the other limitation was we only used biochemical and histological approaches. So, our recommendation for future work is to include different types of approaches for different types of organs of different types of animal models to better understand the exact mechanism (s) of mercury oxide toxicity.

### Acknowledgments

We like to express our gratitude to the Biology Department of the University of Zakho for its support and the resources it provided to us to perform this study.

### Declaration of Conflict of Interest

The authors declare that there is no conflict of interest.

### Funding statement

This study didn't receive any funding support






Fig. 1. Mercury oxide effects on the body weight.

TABLE 1. Effects of mercury oxide on kidney function markers (urea and creatinine) in Wistar rats

Kidney Function Test	Control	Vinegar	0.375 Mg	1.5 Mg	4.5 Mg	P-Value
Urea	43.1± 2.3	45.1 ± 1.8	39.7 ± 4.9	44.86 ± 2.12	45.6 ± 14.1	0.186
Creatinine	0.3 ± 0.01	0.4 ± 0.05	0.4± 0.05	0.4 ± 0.034	0.35 ± 0.03	0.104

**TABLE 2.** Effects of mercury oxide on liver function markers (AST, ALT, ALP, GGT, Albumin, Protein, and LDH) in Wistar rats

Function Liver Tests	Control	Vinegar	0.375 Mg	1.5 Mg	4.5 Mg	P-Value
AST	160.5±68.9	158.6 ± 21.2	215.5±58.1	248.4±49.2 *	123.1±35.3	0.0092**
ALT	56.1 ±21.9	65.1±10.6	63.0±8.2	77.7±25.2	42.9±3.6	0.0476*
ALP	426.4 ±154.4	435.6±147.7	240.0±56.9 *	350.7±103.9	149.0±32.6**	0.0034**
GGT	-1.4 ±1.2	-1.0±1.7	0.0±0.0	0.0±0.0	0.5±0.9	0.1213
Albumin	4.2 ± 1.2	3.9±0.3	4.0±0.1	3.3±0.3*	3.5±0.5	0.0562
Protein	6.7 ±2.1	6.7±0.3	6.6±0.5	6.4±0.3	6.2±0.2	0.2186
LDH	1350.2± 50.6	684.5±337.3	1052.6±505.4	1662.7±498.1	427.8±225.0 *	0.0194*

**Fig. 2.** Transverse section of kidneys showing: (A and B) normal architecture of renal corpuscle and renal tubules in control and vinegar groups respectively. (C and D) showing the breakdown of the glomerulus into two parts (arrow) in groups 4 and 5 respectively (A, B, C, and D 400x).**Fig 3.** Transverse section of the kidney showing: (A and B) shrinkage of glomeruli in group 3 and group 4 respectively. (C) Reveals degeneration of glomeruli and renal tubules (arrow) in group 3. (D) Highlighting extravasation of blood in medulla group 3 (A 100x. B, C and D 400x).




Fig. 4. Illustrates the weights of kidneys (g) in both the control and experimental groups.




Fig. 5. Illustrates the diameter of glomeruli ( $\mu\text{m}$ ) in both the control and experimental groups.




Fig. 6. Transverse section of the liver showing: (A and B) normal histological structures of the hepatocytic plate, hepatic sinusoid, and central vein in control and vinegar groups respectively. (C and D) dilation of a hepatic sinusoid with normal hepatocyte structure in groups 3 and 4 respectively (A, B, C, and D 400x).




Fig. 7. Illustrates the liver weight (g) in both the control and experimental groups.

## References

- Felske A., Vandieken V. and Pauling B.V. Molecular quantification of genes encoding for green-fluorescent proteins. *J. Microbiol. Methods*, **52**, 297–304 (2003).
- Guentzel, J.L., Powell, R.T. and Landing, W.M. Mercury is associated with colloidal material in an estuarine and an open-ocean environment. *Mar. Chem.*, **55**, 177–188 (1996).
- Saleh A.M., Hassan Y.M. and Habeeb T.H. Interactive effects of mercuric oxide nanoparticles and future climate CO<sub>2</sub> on maize plant. *J. Hazard. Mater.*, **40**, 123849 (2021). DOI: 10.1016/j.jhazmat.2020.123849.
- Liu, W., Majumdar, S., Li, W., Keller, A.A. and Slaveykova, V.I. Metabolomics for early detection of stress in freshwater alga *Poteroiochromonas malhamensis* exposed to silver nanoparticles, *Sci. Rep. J.*, **10**, 20563 (2020). DOI: 10.1038/s41598-020-77521-0
- Yasser, M. H., Hamada, A., Ayman, H. Z.Ola Hammouda and Salah-Eldin A. Khodary . Mercuric oxide nanoparticles deferred germination and devastated root anatomy of maize. *Research Square*, **10**(2), 723-735 3579893 (2023). DOI: 10.21203/rs.3.rs-3579893/v1.
- Mohadesi, A., Ranjbar, M. and Hosseinpour-Mashkani, S.M., Solvent-free synthesis of mercury oxide nanoparticles by a simple thermal decomposition method, *Superlattice. Microst.*, **66**, 48–53(2014). DOI: 10.1016/j.spmi.2013.11.017.
- Cassier-Chauvat, C. and Chauvat, F. Responses to oxidative and heavy metal stresses in cyanobacteria: recent advances, *Int. J. Mol. Sci.*, **16**, 871–886(2014). DOI: 10.3390/ijms16010871.
- D. Teng, K. Mao, W. Ali, G. Xu, G. Huang, and N.K. Niazi. Describing the toxicity and sources and the remediation technologies for mercury-contaminated soil, *RSC Adv.* **39** (2020) 23221–23232, DOI: 10.1039/D0RA01507E.
- F. Chen, Z. Xiao, L. Yue, J. Wang, Y. Feng, X. Zhu, Z. Wang, and B. Xing. Algae response to engineered nanoparticles: current understanding, mechanisms and implications, *Environ. Sci. Nano* **6** (2019) 1026–1042, DOI: 10.1039/C8EN01368C.
- Carrie, L. M., David, B. W. and Brian, P. L.. Formation of Soluble Mercury Oxide Coatings: Transformation of Elemental Mercury in Soils. *Environ Sci Technol.*, **49**(20):12105-11(2015).
- Maria, E. C., Gisele, L. M., Susana, I. D. P. and Gabriela, P. F. A. Mercury and human genotoxicity: critical considerations and possible molecular mechanisms. *Pharmacol. Res.*, **60**(4), 212-20(2009). DOI: 10.1016/j.phrs.2009.02.011.
- Mason, R.P., Fitzgerald, W.F., and Morel, F.M.M. The biogeochemical cycling of elemental mercury: Anthropogenic influences. *Geochem. Cosmochim. Acta*, **58**, 3191–3198 (1994).
- Mason, R.P., Rolffhus, K.R., and Fitzgerald, W.F. Mercury in the North Atlantic. *Mar. Chem.*, **61**, 37–53 (1998).
- Lima, L., Bernardez, L.A., and Barbosa, L. Characterization and treatment of artisanal gold mine tailings. *J. Hazard. Mater.*, **150**, 747–753 (2008)
- Sousa, R.N., Veiga, M.M., and Klein, B. Strategies for reducing the environmental impact of reprocessing mercury-contaminated tailings in the artisanal and small-scale gold mining sector: Insights from Tapajos River Basin, Brazil. *J. Clean Prod.*, **18**, 1757–1766 (2010).
- Wang, Q., Kim, D., and Dionysiou, D.D. Sources and remediation for mercury contamination in aquatic systems—A literature review. *Environ. Pollut.*, **131**, 323–336 (2004).
- Ning, P., Guo, X., and Wang, X. Removal of mercury (II), elemental mercury and arsenic from simulated flue gas by ammonium sulphide. *Environ. Technol.*, **36**, 2691–2701(2015).
- Zaki, M. S., Elbattrawy, N., and Fawzi, O. M. Effect of mercuric oxide toxicity on some biochemical parameters in African catfish (*Clarias gariepinus*) present in the River Nile. *Life Science Journal*, **8**(1), 363-368 (2011).
- M. Berlin, R. K. Zalups, and B. A. Fowler, “Mercury,” in *Handbook on the Toxicology of Metals*, G. F. Nordberg, B. A. Fowler, M. Nordberg, and L. T. Friberg, Eds., chapter 33, Elsevier, New York, USA, 3rd edition, 2007.
- M. Nylander, L. Friberg, and B. Lind, “Mercury concentrations in the human brain and kidneys in relation to exposure from dental amalgam fillings,” *Swedish Dental Journal*, vol. **11**, (5), 179–187, (1987).
- L. J. Hahn, R. Kloiber, M. J. Vimy, Y. Takahashi, and F. L. Lorscheider, “Dental ‘silver’ tooth fillings: a source of mercury exposure revealed by whole-body image scan and tissue analysis,” *FASEB Journal*, **3**, (14), 2641–2646 (1989).



22. G. Danscher, P. Horsted-Bindslev, and J. Rungby, "Traces of mercury in organs from primates with amalgam fillings," *Experimental and Molecular Pathology*, **52**, (3), 291–299 (1990).
23. Robin A. Bernhoft. Mercury Toxicity and Treatment: A Review of the Literature. *Journal of Environmental and Public Health*. 2012, DOI:10.1155/2012/460508.
24. Haji, A. A., and Al-Habib, O. A. The Role of K<sup>+</sup>, Ca<sup>2+</sup> channels, some endothelium hyperpolarizing factors in taurine induced vasorelaxation in rat's aorta. *Technium BioChemMed*, **2**(1), 101–109 (2021). DOI: 0.47577/biochemmed.v2i1.3234
25. Cristina Salvador Pedroso, A., Eusébio Reis Gomes, L., and de Carvalho Jorge, M.R. Mercury removal from process sludges via hypochlorite leaching. *Environ. Technol.* **15**, 657–667 (1994).
26. Clarkson, T.W. Mercury: Major issues in environmental health. *Environ. Health Prospect*, **100**, 31–38 (1992).
27. Suresh Kumar Reddy, K., Shoaibi, A.A., and Srinivasakannan, C. Elemental mercury adsorption on sulfur-impregnated porous carbon—A review. *Environ. Technol.* **35**, 18–26 (2014).
28. Miretzky, P., and Cirelli, A.F. Hg (II) removal from water by chitosan and chitosan derivatives: A review. *J. Hazard. Mater.* **167**, 10–23 (2009).
29. Ferreira, G., Santander, A., Chavarría, L., Cardozo, R., Savio, F., Sobrevia, L., and Nicolson, G. L. Functional consequences of lead and mercury exposomes in the heart. *Molecular aspects of medicine*, **87**, 101048 (2022). DOI: 10.1016/j.mam.2021.101048
30. Basu, N., Bastiansz, A., Dórea, J. G., Fujimura, M., Horvat, M., Shroff, E., Weihe, P., and Zastenskaya, I. Our evolved understanding of the human health risks of mercury. *Ambio*, **52**(5), 877–896 (2023). DOI: 10.1007/s13280-023-01831-6
31. Balali-Mood, M., Naseri, K., Tahergorabi, Z., Khazdair, M.R., and Sadeghi, M. Toxic mechanisms of five heavy metals: Mercury, lead, chromium, cadmium, and arsenic. *Front. Pharmacol.* **12**, 643972 (2021).
32. Jayasumana, C., Gunatilake, S. and Siribaddana, S. Simultaneous exposure to multiple heavy metals and glyphosate may contribute to Sri Lankan agricultural nephropathy. *BMC Nephrology* **16**: 103 (2015).
33. Yang, L., Zhang, Y., Wang, F., Luo, Z., Guo, S., and Strähle, U. Toxicity of mercury: Molecular evidence. *Chemosphere*, **245**, 125586 (2020).
34. Renu, K., Chakraborty, R., Myakala, H., Koti, R., Famurewa, A.C., Madhyastha, H., Vellingiri, B., George, A., and Gopalakrishnan, A.V. Molecular mechanism of heavy metals (lead, chromium, arsenic, mercury, nickel and cadmium)-induced hepatotoxicity: A review. *Chemosphere*, **271**, 129735 (2021). DOI: 10.1016/j.chemosphere.2021.129735
35. Rizwan S., Naqshbandi A., Farooqui Z., Khan A. A., and Khan F., Protective effect of dietary flaxseed oil on arsenic-induced nephrotoxicity and oxidative damage in rat kidney, *Food and Chemical Toxicology*, **68**, 99-107, 0278-6915, (2014). DOI: 10.1016/j.fct.2014.03.011
36. Noman, A. S. M., Dilruba, S., Mohanto, N. C., Rahman, L., Khatun, Z., Riad, W. and Haque, A. Arsenic-induced histological alterations in various organs of mice. *Journal of cytology and histology*, **6** (3) (2015).
37. Guzzi, G.P., and La Porta, C.A.M. Molecular mechanisms triggered by mercury. *Toxicology*, **244**, 1–12 (2008). DOI: 10.1016/j.tox.2007.11.002
38. Bridges, C. C., and Zalups, R. K. Mechanisms involved in the transport of mercuric ions in target tissues. *Archives of toxicology*, **91**(1), 63–81 (2017). DOI:10.1007/s00204-016-1803-y

## دراسة في الجسم الحي للتأثيرات الفسيولوجية والنسجية لجرعات مختلفة من أكسيد الزئبق على الكبد والكلية في ذكور جرذان ويستار

عبدالستار عبدالجبار حاجي\*<sup>1</sup>، الان ادريس يوسف<sup>1</sup>، جنار مصطفى محمد<sup>1</sup>، ثامر محمد بشير<sup>1</sup> واحسان حسين محمد<sup>2</sup>.

<sup>1</sup> قسم علوم الأحياء - كلية العلوم - جامعة زاخو- زاخو - إقليم كردستان - العراق.

<sup>2</sup> قسم الفسيولوجيا الطبية والصيدلة - كلية الطب - جامعة دهوك - دهوك - إقليم كردستان - العراق.

### الملخص

يؤثر تراكم الزئبق على الجهاز الهضمي، والكلية. هدفنا في هذه الدراسة هي دراسة التأثيرات الفسيولوجية والنسجية لأكسيد الزئبق على الكبد والكلية في ذكور جرذان ويستار. خلال 22 يوماً، قمنا بتقسيم 25 جرذان إلى 5 مجموعات. أعطيت المجموعة الضابطة الماء فقط. المجموعة التي تناولت الخل فقط أعطيت الخل فقط. أعطيت المجموعة المعالجة بأكسيد الزئبق (0.375 HgO ملجم/كجم/يوم من الزئبق. المجموعة المعالجة بأكسيد الزئبق تعطي 1.5 ملغم/كجم/يوم من الزئبق. أعطيت المجموعة المعالجة بأكسيد الزئبق (4.5 HgO ملجم/كجم/يوم من الزئبق. قمنا بقياس مستويات ALT، AST، LDH، ALP، الألبومين، الكرياتينين، واليوريا. كما تمت دراسة التشريح المرضي للكبد والكلية. لاحظنا تمدد الجيوب الكبدية، و تلف النيبات الكلوية، وانكماش الكبيبة. أظهرت هذه الدراسة اختلافات غير كبيرة بين المجموعات من حيث قطر الكبيبة الكلوية. أظهرت النتائج أن الزئبق بجرعة (1.5 ملغم/كجم/يوم) كان لديه مستويات أعلى بكثير من إنزيمات ALT، LDH، وAST بالمقارنة مع مجموعة السيطرة. أثناء تناول أعلى جرعة من أكسيد الزئبق (4.5 ملجم/كجم/يوم)، انخفضت مستويات إنزيم LDH وALT وAST بالمقارنة مع الجرعات الأقل. أظهرت نتائجنا زيادة غير مهمة في مستوى اليوريا. وبناء على ذلك، أظهرت هذه الدراسة أن تعرض لأكسيد الزئبق يمكن أن يسبب تسمم الكبد والكلية.

**الكلمات الدالة:** أكسيد الزئبق، علم وظائف الأعضاء، الأنسجة، الكبد، الكلية.

\*الباحث الرئيسي: عبدالستار عبدالجبار حاجي. البريد الإلكتروني [Abdulsatar.haji@staff.uoz.edu.krd](mailto:Abdulsatar.haji@staff.uoz.edu.krd)

رقم الموبايل : 07502153874