



Main Compounds, Antioxidant Activity and Lethal Effect of *Thymus pallescens* Essential Oil on Protoscoleces of Hydatid Cysts



Yamina Kadari^{1,2,3}, Sidi Mohammed Ammar Selles^{1,3*}, Mokhtaria Kouidri^{1,2} and Imane Ould Amer^{4,5}

¹ Veterinary Sciences Institute, University of Tiaret, 14000 Tiaret, Algeria.

² Farm animal reproduction laboratory, Veterinary Sciences Institute, University of Tiaret, 14000 Tiaret, Algeria.

³ Laboratory of Research on Local Animal Products, Veterinary Sciences Institute, University of Tiaret, 14000 Tiaret, Algeria.

⁴ Faculty of Life and Nature Sciences, University of Tiaret, Tiaret 14000, Algeria.

⁵ Laboratory of Agro-Biotechnology and Nutrition in Semi-Arid Areas, University of Tiaret, Tiaret 14000, Algeria.

Abstract

THIS work aimed to investigate the chemical constituents, in-vitro scolicidal, and antioxidant activities of *Thymus pallescens*. Hydrodistillation and gas chromatography with flame ionization detection and gas chromatography-mass spectrometry were used for the extract and determination of compounds in essential oil, respectively. The antioxidant activity of essential oil was assessed by DPPH assay and FRAP assay. Collected aseptically from the sheep livers infected with hydatid cysts slaughtered at the Tiaret abattoir, the protoscoleces used in the present study were washed previously with sterile normal saline. Essential oil scolicidal activity was carried out at the dose (4.55 and 9.1 mg/ml) during 5, 10, and 15 min. An average essential oil yield of 1.87 ± 0.30 % (w/w) was obtained in the present study. Carvacrol (57.31%) was found to be the major compound followed by γ -terpinene (14.09 %) and p-cymene (10.64 %). Moderate antioxidant activity was shown for this essential oil with an IC₅₀ value of 1090.03 ± 153.59 μ g/ml and an EC₅₀ value of 1043.51 ± 130.86 μ g/ml. All protoscoleces were killed only after 5 min of exposure to the dose of 9.1 mg/ml. Good scolicidal and moderate antioxidant activities of *Thymus pallescens* essential oil was shown in the present study.

Keywords: Antioxidant activity, Carvacrol, Chemical constituents, Essential oil, Scolicidal activity, *Thymus pallescens*.

Introduction

Considered a neglected and potentially fatal parasitic zoonosis [1], cystic echinococcosis (CE) is an extremely endemic cosmopolitan zoonosis. It is especially prevalent in parts of South America, North Africa, China, and the Middle East [2]. It is caused by the larval stages of the taeniid helminth *Echinococcus granulosus* [3]. Dogs and other canids are definitive hosts for the parasite's adult stage, colonizing their intestines [4]. Infection can occur in intermediate hosts (humans, cattle, sheep, pigs, and horses) through ingestion of infectious eggs shed in dog feces through close contact or environmental

contamination, resulting in the development of CE-containing protoscoleces (larvae stage) in various parts of the body (liver, lungs...etc.) [5, 6]. In humans, the disease poses a health and economic threat (leading to increased medical costs and risk of death) [1, 7]. In animals, this leads to direct and indirect economic losses such as a decrease in milk production, meat quality, wool, skin, and reproduction [7, 8].

According to the American WHO-IWGE classification [9], treatment of cystic echinococcosis depends on several factors; such as the type of cyst, its size, its location, and the presence/absence of

*Corresponding authors: Selles Sidi Mohammed Ammar: E-mail: nuidr10@yahoo.fr and sidimohammedammar.selles@univ-tiaret.dz, Tel: +213-554-200-199; Fax: + 213 46 42 71 (Received 11 October 2024, accepted 08 December 2024)

DOI: 10.21608/EJVS.2025.327474.2421

©2025 National Information and Documentation Center (NIDOC)

complications. The treatment consists of the complete exeresis of the cyst, regardless of its location. If the cyst with all its layers cannot be completely removed, it is necessary to associate protoscolicidal agents to avoid the intraoperative dissemination of fluid rich in protoscoleces during surgery and thus avoid the recurrence of cystic echinococcosis [10].

However, many conventional scolical agents cause undesirable side effects such as sclerosing cholangitis, hepatic necrosis, and methemoglobinemia [11, 12]. The World Health Organization has reported that discovering a new scolical agent that is more effective and produces fewer adverse effects is crucial [13]. Currently, herbs are notable sources of very useful therapeutic agents against many pathological infections [14]. Over the last few decades, many plant species belonging to different families have been used to neutralize the contents of hydatid cysts in vitro, ex-vivo, and in-vivo models [5, 15-18].

One of the most widespread genera in terms of number in the family Lamiaceae is *Thymus* (with more than 350 species growing mainly in the Mediterranean basin and Asia) [19-20]. Among these species, *Thymus palleescens* de Noé (synonym *Thymus fontanesii* Boiss. et Reut.) is a native plant to northern Algeria and is commonly used as a food preservative and in Algerian popular medicine [21]. However, *Thymus* essential oil is a readily accessible natural source of antibiotics, antioxidants, and other medicinal applications, not toxic, and rich in phenolic compounds [22].

Several studies have investigated the scolical activity of various *Thymus* essential oils [23-26]. A previous study carried out by Selles *et al.* [26] determined the chemical composition, scolical, and antioxidant activities of the essential oil of *Thymus fontanesii*. The objectives of the present study were to determine the chemical composition and to confirm the scolical and antioxidant activities of *Thymus palleescens* collected from another region, namely the Sebaine locality (Tiaret region, Algeria).

Material and Methods

Extraction of essential oil

Leaves of *Thymus palleescens* were collected in May 2022 from the Sebaine locality, Tiaret, Algeria (35° 27' 22" N and 1° 36' 13" E). The plant was identified by AIT HAMOU (botanist at the Faculty of Life and Natural Sciences, Ibn Khaldoun University, Tiaret, Algeria), and dried in obscurity at room temperature. A mixture of twenty grams of *Thymus palleescens* dry leaves with 500 ml of distilled water was subjected to hydrodistillation for 3 h at the end of extraction the oily phase was recovered and dried by anhydrous sodium sulfate and stored in a refrigerator at 4 C° in a sealed dark glass

bottle until its use. The extracted essential oil ($1.87 \pm 0.30\%$ (w/w)) was calculated by dividing the weight of essential oil obtained during each hydrodistillation by the weight of dry matter used initially..

Analysis of essential oil

The determination of the chemical composition of the essential oil was carried out by Sarl Pyrenessences Analysis (France), according to the ISO 11024 standard using gas chromatography equipped with a flame ionization detector (GC/FID) and by gas chromatography coupled with mass spectrometry (GC-MS).

Collection of protoscoleces

Sheep livers naturally infected by hydatid cysts were collected from the Tiaret municipal slaughterhouse and transferred to the Parasitology Laboratory (Veterinary Sciences Institute, University Ibn Khaldoun of Tiaret). The cyst fluid was aseptically aspirated by a syringe then transferred into the glass cylinders and allowed to settle for 30 min to facilitate the precipitation of protoscoleces. After removing the supernatant, the protoscoleces were washed three times with sterile normal saline (0.9% NaCl) [27]. Finally, a 0.1% eosin staining test was used to assess the viability of protoscoleces (observation under an optical microscope of muscular movements and the impermeability of the protoscoleces). Live protoscoleces were stored at 4°C in a dark container containing normal saline and for until use.

Scolical activity

Scolical effect of Thymus palleescens

To evaluate the scolical activity of *Thymus palleescens* EO, concentrations of 4.55 and 9.1 mg/ml were previously prepared separately and dissolved in tween 20/ physiological normal saline solution (1/10 v/v) and then mixed well using a vortex mixer (Dragon Lab MX-S) to ensure good dispersion of the essential oils. In each test tube, half ml of protoscoleces solution and half ml of different concentrations of *Thymus palleescens* EO tested were added. The tubes were mixed gently and incubated for 5, 10, and 15 minutes at 37°C, then removing the supernatant [17]. The mortality rate of protoscoleces was studied by adding 0.5 ml of 0.1% eosin and then incubating for 15 min at room temperature. At the end of this coloring, a drop of the mixture was placed on a slide and the percentage mortality of protoscoleces was examined by counting at least 1300 protoscoleces under a light microscope [28]. In addition, physiological normal saline solution and a mixture tween 20/ physiological normal saline solution were used as control groups. Each tests were performed in triplicate.

Viability

The assessment of the muscle movements of protoscolecids under an optical microscope was carried out according to the method described by Miman et al. [29] which consists of exposing for 15 min a mixture of a drop rich in protoscolecids with a drop of 0.1% eosin solution. Afterward, a drop of the mixture is placed on a slide covered with a coverslip. Protoscolecids stained red are considered dead, while colorless ones are alive (figure 1).

Antioxidant Activity

DPPH Free Radical Scavenging Activity

The method previously described by Blois [30] was used to evaluate the free radical scavenging activity of various concentrations *Thymus pallescens* EO tested using 1,1-diphenyl-2-picryl hydrazyl (DPPH). One ml of various concentrations of *Thymus pallescens* EO was added to a 200 µl solution of DPPH free radical in ethanol (0.5 mM). A control group (containing all reagents except the tested essential oil) was also prepared. After that, the mixture was shaken and kept in complete darkness for 30 min. The absorbance was read at 517 nm. The scavenging activity was estimated, by using the equation.

$$\text{DPPH I\%} = \left(\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \right) \times 100$$

Where: A_{control} : Control reaction absorbance;
 A_{sample} : Testing specimen absorbance.

The results are expressed as IC_{50} . All tests were performed in triplicate, and results were expressed as mean values \pm standard deviation (SD). VitC and Gallic acid were used as the positive controls.

Ferric reducing power assay (FRAP)

The determination of the ability of EO to reduce Fe^{3+} to Fe^{2+} was performed according to the method previously described by Vijayalakshmi and Ruckmani [31] with some adjustments. This assay consists of incubation at 50°C for 20 min of a mixture containing an equal volume of various concentrations of *Thymus pallescens* EO, phosphate buffer (0.2 M, pH 6.6), and $C_6N_6FeK_3$ (1%). At the end of this incubation, 0.5 ml of trichloroacetic acid (10%) was added to the mixture and then centrifuged at 3000 rpm for 10 min. After centrifugation, 1 ml of supernatant was taken and mixed with the same volume of distilled water and 0.5 ml of $FeCl_3$ (0.1). The mixture was allowed at room temperature in the dark for 10 min before measuring the absorbance at 700 nm. The results are expressed as EC_{50} . All tests were performed in triplicate, and results were expressed as mean values \pm standard deviation (SD). Ascorbic acid was used as the positive control.

Statistical analysis

One-way analysis of variance (ANOVA) followed by Tukey's highly significant difference

(HSD) post hoc test was performed by R software (version 4.3.1/2023-06-16) to investigate the differences between the test and control groups.

Results

Chemical composition

GC/MS and GC/FID analysis allowed the identification of one hundred and seven compounds, representing 100% of the total components of *Thymus pallescens* EO. Carvacrol was the major compound of this EO (57.31%). Table 1 summarizes the main compounds of *Thymus pallescens* EO.

Antioxidant activity

Regarding the antioxidant activity, table 2 summarizes the IC_{50} and EC_{50} for *Thymus pallescens* EO and standards tested by DPPH free radical scavenging activity and ferric reducing power.

Scolicidal activity

In the present study, *Thymus pallescens* EO exhibited high scolicidal activity. A 100% mortality rate at the doses of 9.1 mg/ml after only 5 min of exposure was observed (figure 2). However, at a concentration of 4.55 mg/ml, this EO killed the protoscolecids with rates of 20.03 %, 31.48 %, and 67.42 % after 5 min, 10 min, and 15 min of exposure, respectively. Additionally, very highly significant ($P < 0.001$) and significant ($P < 0.05$) differences were found between this EO at doses of 9.1 mg/ml and 4.55 mg/ml and controls, respectively.

Discussion

Yield

In the current study, the yield of EO extracted from *Thymus pallescens* dried leaves and produced by simple hydro-distillation ranged from $1.87 \pm 0.30\%$ (w:w). A similar result with a rate of 1.7 to 1.9 % was reported by Benchabane et al. [20] for *Thymus pallescens* harvested from the center of northern Algeria and extracted by Clevenger apparatus at different durations of distillations. Likewise, the rates of *Thymus pallescens* EO collected from Ain Defla (Algeria) by steam distillation, and steam diffusion were $1.54 \pm 0.07\%$ and $1.50 \pm 0.04\%$ respectively [32]. However, the same authors obtained a lower yield ($1 \pm 0.05\%$) using the hydro distillation method. The higher yield was mentioned by Hazzit et al. [19], Hazzit et al. [33], and Selles et al. [26], from *Thymus pallescens* collected from Lakhdaria, Oued Rhiou, and Maalab (Algeria) with a yield ranging from 2.06% to 6.2%.

Yield can be influenced by harvest time and stage of development [19]. These authors found that a maximum yield of 4.6% was obtained at the full flowering stage while a minimum yield of around 0.9% was displayed at the start of the vegetative

cycle. Additionally, the region and harvest period, type, and duration of distillation can influence yield.

Chemical composition

In the present study, carvacrol is the major component of *Thymus pallescens* EO (57.31 %), followed by γ -terpinene (14.09 %), p-cymene (10.64 %), β -myrcene (2.65 %), limonene (2.09 %), α -terpinene (1.92 %), α -pinene (1.40 %), linalool (1.34 %), and α -thujene (1.28%). While this EO contains a small amount of thymol (0.60 %) (Table 1). A similar composition was reported by Selles *et al.* [26] who recorded that carvacrol is the main compound of this essential oil followed by γ -terpinene, p-cymene, α -terpinene, linalool, and α -pinene with slight percentage differences. Similarly, these authors recorded little thymol in this EO (0.49 %). However, low rates of β -myrcene (1.56 %), limonene (0.69 %), and α -thujene (0.36 %) were noted by the same authors for this EO. In the current research, the results are in agreement with those carried out by Moutassem *et al.* [34] which showed that carvacrol was the major component (56.64%) of *Thymus pallescens* EO harvested from various regions of Mascara (Algeria). Similarly, Sadjia *et al.* [32] and Benchabane *et al.* [20] showed that carvacrol was the main chemical constituent with rates respectively of 79.4 - 86.3 % depending on the extraction method used and from 63.3-68.2 % impacted by the distillation time. Likewise, Hazzit *et al.* [19] noted that *Thymus pallescens* EO collected from four regions of Algeria (Sidi Aissa, Boussaada, Kadiria, and El-Asnam) were a carvacrol chemotype (44.4–57.7%) followed by p-cymene (10.3–17.3 %) and γ -terpinene (10.8–14.2 %). Contrariwise, the chemical composition of *Thymus pallescens* EO collected from Oued Rhiou (Algeria) was dominated by thymol (49.3 %), while carvacrol only represents a small amount (9.0%) [19].

Figueiredo *et al.* [35] noted that the geographical location of *Thymus pallescens* is Algeria and Tunisia and that the chemical composition of its essential oil is as follows: Thymol >1-68 %, Carvacrol 9-65 %, γ -terpinene 6-17 %, p-cymene 5-17%, Linalool >1-7 %, α pinene >1-6%.

The difference in *Thymus pallescens* EO composition can be explained by hydrodistillation time [20], method of extraction [36], and part of the plant [19]. In addition, geographic location, season, and moment of harvesting plant affect the composition of essential oil [37-38].

Antioxidant activity

Reducing power and DPPH assay were used to determine the ability of EO to reduce ferric ions (Fe³⁺) to ferrous ions (Fe²⁺) and the capacity to scavenge the stable DPPH•. IC₅₀ and EC₅₀ values of reducing power and DPPH scavenging activity of

Thymus pallescens EO and standards were summarized in Table 2.

This study showed an IC₅₀ of *Thymus pallescens* EO of 1090.03 ± 153.59 µg/ml. Higher results were cited by Ziani *et al.* [39], Benchabane *et al.* [20], and Hazzit *et al.* [40] with an IC₅₀ of 103g/ml, 149.8 mg/ml, and 410.2 mg/ml, respectively. Several studies have reported low IC₅₀ of the essential oil of specimens of *Thymys fontanesii* harvested from various regions of Algeria ranging from 15.58 µg/ml to 346.08 µg/ml [26, 41-42].

In the current investigation, less IC₅₀ was recorded by ascorbic acid (4.45 µg/ml) and gallic acid (3.19 µg/ml). A very highly significant difference (P=0.00000752) between standard molecules and *Thymus pallescens* EO was demonstrated.

However, the comparison of the result of the present research with other species of *Thymus* shows a powerful activity of *Thymus vulgaris* with an IC₅₀ of 0.19 µg/ml and 0.5 µg/ml, respectively [43-44]. This potent DPPH free radical scavenging activity can be explained by its richness in thymol [45]. Likewise, a strong IC₅₀ (0.08 mg/ml) was shown by Zaïri *et al.* [22] for the *Thymus capitatus* EO carvacrol chemotype. Additionally, He *et al.* [38] reported a low IC₅₀ of *Thymus quinquecostatus celak* collected from different regions of China with a value ranging from 0.512 mg/ml to 0.931 mg/ml.

Our finding recorded higher EC₅₀ (1043.51 µg/ml) for *Thymus pallescens* EO comparatively to ascorbic acid (EC₅₀ = 23.39 µg/ml). A lower result was noted by Selles *et al.* [26] with a value of 869.82 µg/ml *Thymus fontanesii* EO. Likewise, low EC₅₀ values (63.8 µg/ml to 112.6 µg/ml) of various samples of *Thymus fontanessi* EO were reported by Sid Ali *et al.* [41]. A higher concentration was found in a study realized by Ziani *et al.* [39] with an EC₅₀ of 63 mg/ml. A highly significant difference (P=0.00187) between standard molecules and *Thymus pallescens* EO was demonstrated.

The high content of this essential oil in phenolic components (carvacrol) explains this good antioxidant activity. Moreover, hydro distillation time [20], method and the solvents used [46], geographic location, radiation exposure [21], the chromatographic profile of the putative antioxidant, and its polarity also have consequences on the antioxidant properties [38, 43].

Scolicidal activity

In our investigation, *Thymus pallescens* EO exhibited strong scolicidal activity at very low concentrations and in a short time. A lethality of 100% at 5 min of exposure was recorded by this essential oil (figure 2). While the dose of 4.55 mg/ml

killed 20.03%, 31.48%, and 67.42% after 5 min, 10 min and 15 min of exposure, respectively.

Selles et al. [26] noted that the essential oil of *Thymus fontanesii* possesses a powerful scolicidal activity. Since this EO at the concentration of 9.25 mg/ml kills 100% of protoscoleces after only 5min of exposure and showed a significant difference with the controls ($P < 0.05$). The same authors noticed no significant difference in scolicidal activity at the concentration of 4.625 mg/ml and the controls. Likewise, Hizem et al. [25] recorded that *thymus capitatus* EO exerts a protoscolicidal effect of 100% at doses of 2 and 3 mg/ml after 5 min and 1 min, respectively. Several previous studies report the potent scolicidal activity of various essential oils with carvacrol major compound depending on dose and time. Mahmoudvand et al. [47] mentioned that 10 mg/ml *Nigella sativa* EO kills 100% of protoscoleces after 10 min of exposure. Similarly, at a concentration of 0.05% *Origanum minutiflorum* EO causes mortality of all protoscoleces after 5 min of exposure [48]. Whereas, Soleimani et al. [49] report that *Origanum vulgare* EO at a concentration of 1% requires 15 min to kill all protoscoleces. Likewise, Moazeini et al. [50] found that 1mg/ml of *Satureja hortensis* EO kills 100% of protoscoleces after 20 min. However, 2 mg/ml of these EOs need 10 min to kill all protoscoleces. Nevertheless, at a dose of more than 17µg/ml, *Zataria multiflora* EO kills protoscoleces after 10 min of exposure [51].

Additionally, previous investigations have shown a powerful scolicidal activity of various EOs but with a longer action time and/or a higher concentration than that observed in the present study [5, 15-16,18, 52-54].

Our study was characterized by low concentration and short time of acting. These criteria are in agreement with the main criteria for an ideal scolicidal agent reported by Moazeni and Larki [55]. This strong scolicidal activity of *Thymus pallescens* EO is due probably to the richness of this essential oil in carvacrol. Previous studies reported that carvacrol has a strong effect against protoscoleces this activity may be linked to the hydroxyl group, which may act as proton translocators [56].

TABLE 1. Main compounds of *Thymus pallescens* EO

| Compounds | % |
|------------------|--------------|
| Carvacrol | 57.31 |
| γ-terpinene | 14.09 |
| p-cymene | 10.64 |
| β-myrcene | 2.65 |
| Limonene | 2.09 |
| α-terpinene | 1.92 |
| α-pinene | 1.40 |
| Linalool | 1.34 |
| α-thujene | 1.28 |
| Thymol | 0.60 |

Furthermore, this experiment allowed us to observe morphological changes in dead protoscoleces such as loss of hooks, contraction of soma, and modification in the shape (square shape) (figure 3). Similar morphological changes were shown by Selles et al. [26]. These authors assume that carvacrol (the major compound of this essential oil) is responsible for these modifications.

Conclusion

In conclusion, carvacrol is the major compound of *Thymus pallescens* essential oil that has shown good potential as an antioxidant and scolicide. Based on our experimental results, we can suppose that this oil could be used as an alternative scolicide to routinely used chemicals during surgical excision of hydatid cysts. However, further ex vivo and in vivo studies are needed to confirm the use of this essential oil as a scolicidal agent during hydatid cyst surgery.

Acknowledgments

The authors would like to thank all veterinary inspectors of the Tiaret slaughterhouse, and the staff of the parasitology laboratory of the Institute of Veterinary Sciences for their help and collaboration. The authors are also grateful to Mr. Ait Hammou Mohamed (Botanist) from the Faculty of Natural and Life Sciences of the University of Tiaret for their collaboration during this study.

Funding statement

No funding was allocated to this study.

Declaration of competing interest

The authors have no conflicts of interest in this study.

Ethical of approval

The samples were collected according to the protocol approved by the scientific committee of the Institute of Veterinary Sciences, Ibn Khaldoun University, Tiaret, Algeria (591/ Vice Rectorate of Third Cycle Higher Education and University Accreditation, Scientific Research and Postgraduate Higher Education / 2021).

TABLE 2. Reducing power and DPPH assays of EO

| | Reducing power (EC ₅₀) µg/ml | DPPH (IC ₅₀) µg/ml |
|-----------------------------|---|-----------------------------------|
| Gallic acid | - | 3.19 ± 0.41 ^a |
| Ascorbic acid | 23.39 ± 0.81 ^a | 4.45 ± 1.05 ^a |
| <i>Thymus pallescens</i> EO | 1043.51 ± 130.86 ^b | 1090.03 ± 153.59 ^b |

Values are expressed as mean ± standard deviation (n = 3). Values with different letters are statistically different at P < 0.05 as measured by one-way analysis of variance (ANOVA)




Fig. 1. A: Live protoscolecids; B: Dead protoscolecids of hydatid cysts after staining with 0.1% eosin GX40




Fig. 2. Scolical activity of *Thymus pallescens* EO against protoscolecids.




Fig. 3. Some modifications of the protoscolex. A: Hook release. B: Change in the shape of the protoscolex (Squar shape) GX40.

References

- Jalil, P.J., Shnawa, B.H. and Al-Ali, S.J., Immunological aspects of cystic echinococcosis: an overview. In: Abbas, R.Z., Khan, A., Liu P., and Saleemi M.K., editor Animal Health Perspectives. Unique Scientific Publishers, Faisalabad, Pakistan: Vol. **I**, pp. 103-113, (2022). <https://doi.org/10.47278/book.ahp/2022.14>
- Harandi, M.F., Budke, C.M. and Rostami, S., The monetary burden of cystic echinococcosis in Iran. *PLoS Negl. Trop. Dis.*, **6**, e1915 (2012). <https://doi.org/10.1371/journal.pntd.0001915>.
- Ali, R., Khan, S., Khan, M., Adnan, M., Ali, I., Khan, T.A., Haleem, S., Rooman, M., Norin, S. and Khan, S.N., A systematic review of medicinal plants used against *Echinococcus granulosus*. *PLoS One.*, **15** (10), e0240456 (2020). <https://doi.org/10.1371/journal.pone.0240456>.
- Hassanzadeh, E., Khademvatan, S., Jafari, B., Jafari, A. and Yousefi, E., In vitro and in silico scolical effect of sanguinarine on the hydatid cyst protoscolices. *PLoS One.*, **18** (10), e0290947 (2023). <https://doi.org/10.1371/journal.pone.0290947>.
- Moazeni, M., Saharkhiz, M.J. and Hosseini, A.A., In vitro lethal effect of ajowan (*Trachyspermum ammi* L.) essential oil on hydatid cyst protoscolices. *Vet. Parasitol.*, **187**(1-2), 203-208 (2012a). <https://doi.org/10.1016/j.vetpar.2011.12.025>
- Mahmoudvand, H., Asadi, A., Harandi, M.F., Sharififar, F., Jahanbakhsh, S. and Dezaki, E.S., In vitro lethal effects of various extracts of *Nigella sativa* seed on hydatid cyst protoscolices. *Iran. J. Basic. Med. Sci.*, **17**(12), 1001 (2014).
- Moghadaszadeh, M., Khayyati, M., Spotin, A., Norouzi, R., Pagheh, A.S., Oliveira, S.M.R., Pereira, M.L. and Ahmadpour, E., Scolicidal and apoptotic activities of 5-hydroxy-1, 4-naphthoquinone as a potent agent against *Echinococcus granulosus* protoscolices. *Pharmaceuticals*. (Basel), **14** (7), 623 (2021). <https://doi.org/10.3390/ph14070623>.
- Garedaghi, Y., Bouree, P., Khaki, A., Shaarba, F. and Rangraz, M.H.A., Evaluation of the effects of Bunium persicum and albendazole on killing hydatid cyst protoscolices in vitro. *Crescent. J. Med. Biol. Sci.*, **9** (2), 89-92 (2022). <https://doi.org/10.34172/cjmb.2022.15>.
- McManus, D.P., Gray, D.J., Zhang, W. and Yang, Y., Diagnosis, treatment, and management of Echinococcosis. *B. M. J.*, **344**, e3866 (2012). <https://doi.org/10.1136/bmj.e3866>.
- Brunetti, E., Kern P. and Vuitton, D.A., Expert consensus for the diagnosis and treatment of cystic and alveolar echinococcosis in humans. *Acta. Trop.*, **114**, 1-16 (2010). <https://doi.org/10.1016/j.actatropica.2009.11.001>.
- Rajabi, M.A., Fatal reactions and methaemoglobinaemia after silver nitrate irrigation of hydatid cyst. *Surg. Prac.*, **13**, 2-7 (2009). <https://doi.org/10.1111/j.1744-1633.2008.00427.x>
- Gholami, S.H., Rahimi-Esboei, B., Ebrahimzadeh, M.A. and Pourhajibagher, M., In vitro effect of *Sambucus ebulus* on scolices of Hydatid cysts. *Eur. Rev. Med. Pharmacol. Sci.*, **17**(13), 1760-1765 (2013).
- Kheirandish, F., Marzban, A., Masoori, L., Nourmohammadi, M., Moradpour, K. and Cheraghpour, K., Efficacy of thymol on the scolices of hydatid cyst. *Herb. Med. J.*, **5** (1), 37-44 (2020). <https://doi.org/10.22087/hmj.v5i1.786>.
- Houshmand, E., Kamalifar, H. and Elmi, H., In vitro Scolicidal Effect of Ginger (*Zingiber officinale* Roscoe) Ethanolic Extract Against Protoscolices of Hydatid Cyst. *Iran. J. Vet. Med.*, **13**(1), 87-99 (2019). <https://doi.org/10.22059/ijvm.2018.263561.1004915>
- Moazeni, M., Saharkhiz, M.J., Hoseini, A.A. and Alavi, A.M., In vitro scolical effect of *Satureja khuzistanica* (Jamzad) essential oil. *Asian. Pac. J. Trop. Biomed.*, **2** (8), 616-620 (2012b). [https://doi.org/10.1016/S2221-1691\(12\)60107-5](https://doi.org/10.1016/S2221-1691(12)60107-5).
- Mahmoudvand, H., Fallahi, S., Mahmoudvand, H., Shakibaie, M., Harandi, M.F. and Dezaki, E.S., Efficacy of *Myrtus communis* L. to inactivate the hydatid cyst protoscolices. *J. Investig. Surg.*, **29** (3), 137-143 (2016). <https://doi.org/10.3109/08941939.2015.1088601>.

17. Selles, S.M.A., Kouidri, M., Belhamiti, T.B., Ait Amrane, A., Benahmed, M. and Hachemi, A., Correction to: Main compounds and in vitro effectiveness of *Syzygium aromaticum* essential oil on protoscoleces of hydatid cyst. *Comp. Clin. Pathol.*, **29**, 911 (2020). <https://doi.org/10.1007/s00580-020-03133-9>
18. Alyousif, M.S., Al-Abodi, H.R., Almohammed, H., Alanazi, A.D., Mahmoudvand, H., Shalamzari, M.H. and Salimikia, I., Chemical composition, apoptotic activity, and antiparasitic effects of *Ferula macrecolea* essential oil against *Echinococcus granulosus* protoscoleces. *Molecules*, **26** (4), 888 (2021). <https://doi.org/10.3390/molecules26040888>
19. Hazzit, M., Baaliouamer, B., Verissimo, A.R., Faleiro, M.L. and Miguel, M.G., Chemical composition and biological activities of Algerian *Thymus* oils. *Food. Chem.*, **116**, 714–721 (2009). <https://doi.org/10.1016/j.foodchem.2009.03.018>
20. Benchabane, O., Hazzit, M., Mouhouche, F. and Baaliouamer, A., Influence of extraction duration on the chemical composition and biological activities of essential oil of *Thymus pallescens* de Noé. *Arab. J. Sci. Eng.*, **40**, 1855-1865 (2015). <https://doi.org/10.1007/s13369-015-1694-x>
21. Alloun, K., Benchabane, O., Hazzit, M., Mouhouche, F., Baaliouamer, A., Chikhoun, A. and Benchabane, A., Effect of gamma ray irradiation on chemical composition, Antioxidant, antimicrobial, and insecticidal activities of *Thymus pallescens* essential oil. *Acta. Chromatogr.*, **31** (1), 57-62 (2019). <https://doi.org/10.1556/1326.2017.00346>
22. Zaïri, A., Nour, S., Zarrouk, A., Haddad, H., Khélifa, A., Achour, L., Tangy, F., Chaouachi, M. and Trabelsi, M., Chemical composition, Fatty acids profile and biological properties of *Thymus capitatus* (L.) Hoffmanns, essential Oil. *Sci. Rep.*, **9** (1), 20134 (2019). <https://doi.org/10.1038/s41598-019-56580-y>
23. Yones, D.A., Taher, G.A. and Ibraheim, Z.Z., In vitro effects of some herbs used in Egyptian traditional medicine on viability of protoscolices of hydatid cysts. *Korean. J. Parasitol.*, **49** (3), 255 (2011). <https://doi.org/10.3347/kjp.2011.49.3.255>
24. Pensel, P., Maggiore, M., Gende, L., Eguaras, M., Denegri, M. and Elisondo, M., Efficacy of essential oils of *Thymus vulgaris* and *Origanum vulgare* on *Echinococcus granulosus*. *Interdiscip. Perspect. Infect. Dis.*, **2014**, 693289 (2014). <https://doi.org/10.1155/2014/693289>
25. Hizem, A., M'rad, S., Oudni-M'rad, M., Mezhoud, H., Ben Jannet, H. and Flamini, G., In-vitro scolicidal activity of *Thymus capitatus* Hoff. Et Link. essential oil on *Echinococcus granulosus* protoscoleces. *J. Essent. Oil. Res.*, **32** (2), 178–185 (2020). <https://doi.org/10.1080/10412905.2019.1711212>
26. Selles, S.M.A., Belhamiti, B.T., Kouidri, M., Ait Amrane, A., Kadari, Y., Kaddour, Z. and Kebrit, S., Chemical compounds, antioxidant and scolicidal potencies of *Thymus fontanesii* essential oil. *Exp. Parasitol.*, **257**, 108699 (2024). <https://doi.org/10.1016/j.exppara.2024.108699>
27. Moazeni, M., Borji, H., Darbandi, M.S. and Saharkhiz, M.J., In vitro and in vivo antihydatid activity of a nano emulsion of *Zataria multiflora* essential oil. *Res. Vet. Sci.*, **114**, 308-312 (2017). <https://doi.org/10.1016/j.rvsc.2017.06.003>
28. Lashkarizadeh, M.R., Asgaripour, K., Dezaki, E.S. and Harandi, M.F., Comparison of scolicidal effects of amphotricin B, silver nanoparticles, and *Foeniculum vulgare* Mill on hydatid cysts protoscoleces. *Iran. J. Parasitol.*, **10** (2), 206–212 (2015).
29. Miman, O., Aycan, O.M., Aydin, C. and Atambay, M., What should be the concentration of eosin to qualification of ideal staining for viability determination on hydatid cyst? *Turk. Hij. Den. Biyol. Derg.*, **67** (1), 21–26 (2010).
30. Blois, M.S., Antioxidant determinations by the use of a stable free radical. *Nature.*, **181**(4617), 1199-1200 (1958). <https://doi.org/10.1038/1811199a0>
31. Vijayalakshmi, M. and Ruckmani, K., Ferric reducing anti-oxidant power assay in plant extract. *Bangladesh. J. Pharmacol.*, **11** (3), 570–572 (2016). <https://doi.org/10.3329/bjpv.v11i3.27663>
32. Sadjia, B., Naima, S. and Chahrazed, B., Extraction of thyme (*Thymus pallescens* de Noé) essential oil by steam-distillation, steam-diffusion and hydro distillation processes: optimization of operating conditions and antioxidant activity. *J. Essent. Oil-Bear. Plants.*, **15** (2), 336-347 (2012). <https://doi.org/10.1080/0972060X.2012.10644056>
33. Hazzit, M., Baaliouamer, A. and Douar-Latreche, S., Effect of heat treatment on the chemical composition and the antioxidant activity of essential oil of *Thymus pallescens* de Noé from Algeria. *J. Essent. Oil. Res.*, **25** (4), 308-314 (2013). <https://doi.org/10.1080/10412905.2013.774626>
34. Moutassem, D., Bellik, Y. and Sannef, M.E.H., Toxicity and repellent activities of *Thymus pallescens* and *Cymbopogon citratus* essential oils against *Sitophilus granaries*. *Plant. Prot. Sci.*, **57** (4), 297-309 (2021). <https://doi.org/10.17221/185/2020-PPS>
35. Figueiredo, A.C., Barroso, J.G. and Pedro, L.G., Volatiles from *Thymbra* and *Thymus* species of the western Mediterranean basin, Portugal and Macaronesia. *Nat. Prod. Commun.*, **5** (9), 1465-1476 (2010). <https://doi.org/10.1177/1934578X10005009>
36. Caputo, L., Amato, G., de Bartolomeis, P., De Martino, L., Manna, F., Nazzaro, F., De Feo, V. and Barba, A. A., Impact of drying methods on the yield and chemistry of *Origanum vulgare* L. essential oil. *Sci. Rep.*, **12** (1), 3845 (2022). <https://doi.org/10.1038/s41598-022-07841-w>
37. Jerković, I., Mastelić, J. and Miloš, M., The impact of both the season of collection and drying on the volatile constituents of *Origanum vulgare* L. ssp. *hirtum* grown wild in Croatia. *International Journal of Food Science & Technology.*, **36** (6), 649-654 (2001). <https://doi.org/10.1046/j.1365-2621.2001.00502.x>
38. He, T., Li, X., Wang, X., Xu, X., Yan, X., Li, X., Sun, S., Dong, Y., Ren, X., Liu, X., Wang, Y., Sui, H., Xia, Q. and She, G., Chemical composition and anti-oxidant potential on essential oils of *Thymus quinquecostatus*

- Celak. from Loess Plateau in China, regulating Nrf2/Keap1 signaling pathway in zebrafish. *Sci. Rep.*, **10** (1), 11280 (2020). <https://doi.org/10.1038/s41598-020-68188-8>.
39. Ziani, B.E., Calhelha, R.C., Barreira, J.C., Barros, L., Hazzit, M. and Ferreira, I.C., Bioactive properties of medicinal plants from the Algerian flora: Selecting the species with the highest potential in view of application purposes. *Ind. Crops. Prod.*, **77**, 582-589 (2015). <https://doi.org/10.1016/j.indcrop.2015.09.034>
40. Hazzit, M., Baaliouamer, A., Faleiro, M.L. and Miguel, M.G., Composition of the essential oils of Thymus and Origanum species from Algeria and their antioxidant and antimicrobial activities. *J. Agric. Food. Chem.*, **54** (17), 6314-21 (2006). <https://doi.org/10.1021/jf0606104>
41. Sid Ali, L., Brada, M., Fauconnier, M-L. and Lognay, G., Chemical Composition and Antioxidant Activity of Thymus fontanesii Essential Oil from Algeria. *The Natural Products Journal.*, **8**, 1-7 (2018). <https://doi.org/10.2174/2210315508666180427162542>
42. Mebarki, N., Nabiev, M., Ziane, H., Chader, H. and Fazouan, F., Evaluation of biological activities of Thyme extracts and their use as a natural alternative to antibiotics for the reduction of pharmaceutical waste. *Algerian Journal of Environmental Science and Technology.*, **6** (1), 1272-1283 (2020).
43. Bozin, B., Mimica-Dukic, N., Simin N. and Anackov, G., Characterization of the volatile composition of essential oils of some Lamiaceae spices and the antimicrobial and antioxidant activities of the entire oils. *J. Agric. Food. Chem.*, **54** (5), 1822-1828 (2006). <https://doi.org/10.1021/jf051922u>
44. Alibi, S., Selma, W.B., Ramos-Vivas, J., Smach, M.A., Touati, R., Boukadida, J., Navas, J. and Ben Mansour, H., Anti-oxidant, antibacterial, anti-biofilm, and anti-quorum sensing activities of four essential oils against multidrug-resistant bacterial clinical isolates. *Curr. Res. Transl. Med.*, **68** (2), 59-66 (2020). <https://doi.org/10.1016/j.retram.2020.01.001>
45. Nezhadali, A., Nabavi, M., Rajabian, M., Akbarpour, M., Pourali, P. and Amini, F., Chemical variation of leaf essential oil at different stages of plant growth and in vitro antibacterial activity of Thymus vulgaris Lamiaceae, from Iran. *Beni-Suef University Journal of Basic and Applied Sciences*, **3** (2), 87-92 (2014). <https://doi.org/10.1016/j.bjbas.2014.05.001>
46. Pérez-Jiménez, J. and Saura-Calixto, F., Effect of solvent and certain food constituents on different antioxidant capacity assays. *Food. Research. International.*, **39** (7), 791-800 (2006). <https://doi.org/10.1016/j.foodres.2006.02.003>
47. Mahmoudvand, H., Dezaki, E.S., Kheirandish, F., Ezatpour, B., Jahanbakhsh, S. and Harandi, M.F., Scolicidal effects of black cumin seed (*Nigella sativa*) essential oil on hydatid cysts. *Korean. J. Parasitol.*, **52** (6), 653-659 (2014). <https://doi.org/10.3347/kjp.2014.52.6.653>.
48. Sokmen, A., Abdel-Baki, A.A.S., Al-Malki, E.S., Al-Quraishy, S. and Abdel-Haleem, H.M., Constituents of essential oil of *Origanum minutiflorum* and its in vitro antioxidant, scolicidal and anticancer activities. *Journal of King Saud University-Science.*, **32** (4), 2377-2382 (2020). <https://doi.org/10.1016/j.jksus.2020.03.018>
49. Soleimani, E., Sarmadian, H., Arjomand Zadegan, M., Ghasemikhah, R. and Ahmadi, H.T., Toxicity and antiparasitic efficacy of essential oils: Analyses of the biochemical compositions and potencies. *Iran. J. Toxicol.*, **15** (1), 1-8 (2021). <http://dx.doi.org/10.32598/ijt.15.1.727.1>
50. Moazeni, M., Saharkhiz, M.J. and Alavi, A.M., The lethal effect of a nano emulsion of *Satureja hortensis* essential oil on protoscolices and germinal layer of hydatid cysts. *Iran. J. Parasitol.*, **14** (2), 214 (2019).
51. Kavooosi, G., and Purfard, A.M., Scolicidal effectiveness of essential oil from *Zataria multiflora* and *Ferula assafoetida*: disparity between phenolic monoterpenes and disulphide compounds. *Comp. Clin. Pathol.*, **22**, 999-1005 (2013). <https://doi.org/10.1007/s00580-012-1518-2>.
52. Mahmoudvand, H., Mirbadie, S.R., Sadooghian, S., Harandi, M.F., Jahanbakhsh, S. and Dezaki, E.S., Chemical composition and scolicidal activity of *Zataria multiflora* Boiss essential oil. *J. Essent. Oil. Res.*, **29** (1), 42-47 (2017). <https://doi.org/10.1080/10412905.2016.1201546>.
53. Tabari, M.A., Youssefi, M.R., Nasiri, M., Hamidi, M., Kiani, K., Alian Samakkhah, S. and Maggi, F., Towards green drugs against cestodes: Effectiveness of *Pelargonium roseum* and *Ferula gummosa* essential oils and their main component on *Echinococcus granulosus* protoscolices. *Vet. Parasitol.*, **266**, 84-87 (2019). <https://doi.org/10.1016/j.vetpar.2018.12.019>
54. Kozan, E., Ilhan, M., Tümen, I. and Küpeli Akkol, E., The scolicidal activity of the essential oil obtained from the needles of *Pinus nigra* Arn. subsp. *pallasiana* (Lamb.) Holmboe on hydatid cyst. *J. Ethnopharmacol.*, **235**, 243-247 (2019). <https://doi.org/10.1016/j.jep.2019.02.018>.
55. Moazeni, M. and Larki, S., In vitro effectiveness of acidic and alkaline solutions on scolices of hydatid cyst. *Parasitol. Res.*, **106**, 853-856 (2010). <https://doi.org/10.1007/s00436-010-1723-3>.
56. Fabbri, J., Maggiore, M.A., Pensel, P. E., Denegri, G. M., Gende, L.B. and Elissondo, M.C., In vitro and in vivo efficacy of carvacrol against *Echinococcus granulosus*. *Acta. Tropica.*, **164**, 272-279 (2016). <https://doi.org/10.1016/j.actatropica.2016.09.001>

المركبات الرئيسية ونشاط مضادات الأكسدة والتأثير القاتل لزيت الأساسي للزعتر على الرؤيسات الأولية للكيسات المائية

قداري يمينة^{1,2,3}، سلس سيدي محمد عمار^{1,3*}، قويدري مختار^{1,2}، ولد عامر إيمان^{4,5}

¹ معهد العلوم البيطرية، جامعة تيارت، تيارت 14000، الجزائر.

² مختبر تكاثر حيوانات المزرعة - جامعة تيارت - الجزائر.

³ مختبر البحث في المنتجات الحيوانية المحلية، جامعة تيارت، تيارت 14000، الجزائر.

⁴ كلية علوم الحياة والطبيعة، جامعة تيارت، تيارت 14000، الجزائر.

⁵ مختبر التكنولوجيا الحيوية الزراعية والتغذية في المناطق شبه الجافة، جامعة تيارت، تيارت 14000، الجزائر.

الملخص

كان الهدف من هذا البحث هو دراسة المكونات الكيميائية، التأثير القاتل لزيت الأساسي للزعتر على حيوية الرؤيسات الأولية للمشوكة الحبيبية للكيس العذري (دراسة في الزجاج) وكذا أنشطة مضادات الأكسدة. تم استخراج الزيت الأساسي عن طريق التقطير المائي وتحليله بواسطة كروماتوغرافيا الغاز المدمج مع المطيافية الكتلية و كروماتوغرافيا الغاز المدمج مع كاشف التآين باللهب. تم استخدام طريقة DPPH و القوة الاختزالية لتقييم نشاط مضادات الأكسدة بواسطة. تم جمع الرؤيسات الأولية للمشوكة الحبيبية للكيس العذري المستخدمة في الدراسة الحالية بطريقة معقمة من أكباد الأغنام المصابة بأكياس الكيس المائي والتي تم ذبحها في مسلخ تيارت، وتم غسلها مسبقاً بمحلول ملحي عادي معقم. تم تقييم النشاط القاتل لزيت الأساسي للزعتر على حيوية الرؤيسات الأولية من خلال تعريض رواسب البروتوسوكولس لتراكيزين من الزيت العطري (4.55، 9.1 ملجم / مل) وضمن جدول زمني لمدة 5، 10 و 15 دقيقة. بلغ مردود استخلاص الزيت الأساسي $1.87 \pm 0.30\%$ (وزن/وزن). كان الكارفاكروول (57.31%) هو المركب الرئيسي للزيت الأساسي الذي تمت دراسته. أظهر نشاط مضادات الأكسدة بواسطة مقايسة DPPH قيمة IC_{50} تبلغ 1090.03 ± 153.59 ميكروغرام/مل، بينما أعطى اختبار قدرة الاختزال الحديدي قيمة EC_{50} تبلغ 1043.51 ± 130.86 ميكروغرام/مل. تم قتل جميع الخلايا الأولية بعد 5 دقائق فقط من التعرض لجرعة 9.1 ملغ/مل. وقد أظهرت الدراسة الحالية أن زيت الزعتر العطري له نشاط جيد في قتل الجراثيم ومضادات الأكسدة المعتدلة.

الكلمات الدالة: المكونات الكيميائية، النشاط القاتل للرؤيسات الأولية، النشاط المضاد للأكسدة، الزيت الأساسي، الزعتر، الكارفاكروول.