



A Trial for Improvement of Kareish Cheese Quality by Using Chitosan Nanoparticles

Rasha M.H. Sayed-Elahl¹, Noha, M. El-Shinawy² and Khaled Nagy³

¹Department of Mycology and Mycotoxins, Animal Health Research Institute (AHRI), Agriculture Research Centre, Cairo, Egypt.

²Food Safety Reference Laboratory, (AHRI), Agriculture Research Centre, Cairo, Egypt.

³Department of Food engineering, packaging and packaging engineering. Agriculture Research Centre, Cairo, Egypt.



CrossMark

THE PRESENT study was undertaken to evaluate the addition of nano-chitosan to kareish cheese for improvement their characters, including odour, tastes, colors and mycological quality. Out of 40 samples of kareish cheese were obtained and mly from retail farmers and supermarkets from different district in Giza Governorate, all samples were contaminated with fungi; the mean count/g was $(1 \times 10^3 \pm 2.4)$. The most common isolated mould species were *Aspergillus* spp. which isolated from (80%) meanwhile; all samples were contaminated with different yeast species. The residues of AFM₁ were detected in 42 %, and the accurate quantitative detection was observed in immunoaffinity column method than thin layer chromatography. The results revealed that the addition or coating with nano-chitosan (0.25% and 0.5%) before or after manufacturing of kareish cheese could cause their prolonged safe preservation as the nano-chitosan have the antimicrobial potential against several bacteria and fungi. Also, the colour, odour and taste of coated or added CSNPs cheeses were more accepted and palatable good characters were observed for a longer time of storage. Hence, it is anticipated that chitosan nanoparticles have the potential of becoming a powerful and safe natural antifungal agent. Therefore, the present study may provide the possibility of the addition of CSNPs to coating kareish cheese manufacture for quality improvement and extending its shelf life.

Keywords: Milk, Kareish cheese, AFM₁, Chitosan nanoparticles, Coating, Antifungal.

Introduction

The contamination of milk and its dairy products with different species of fungi and mycotoxins can be occurred very widely constituting a public health hazard due to the progressive increase in the rates of consumption of these foods [1]. The fungal contamination of milk and milk products including mycotoxigenic fungi which produce mycotoxins that have health hazard for human. The growth of moulds is common and leads to change in colour, odour and flavour; consequently, contamination of

these products leads to significant economic loss and this product become unfit for human consumption [2].

The global problem of mycotoxicosis, particularly in humid regions, including African countries are continuous of significant health hazard. Some moulds are capable of producing more than one mycotoxin, and more than one fungal species are able to produce some mycotoxins. Where as, they have dangerous effects upon consumers is ranging from acutely toxic to immuno suppressive or carcinogenic

*Corresponding author: Rasha M. Sayed-Elahl, E-mail: rasha_hamza2005@hotmail.com.

(Received 21/11/2019, accepted 22/01/2020)

DOI.10.21608/ejvs.2020.19943.1135

©2019 National Information and Documentation Center (NIDOC)

impact to the liver [3,4]. Currently, they are hepatotoxic, nephrotoxic and immuno suppression for animals [5]. Mean while, reduced feed intake, decreased feed utilization, loss in weight gain, reduced reproductive capacity and even death may occur as a result of mycotoxicosis and leading to severe economic injuries [6]. Mycotoxins can cause variety of short term as well as long-term health effects, ranging from immediate toxic responses to potential long-term carcinogenic and teratogenic effects [7,8].

On the other hand, milk is the most animal product exposed to mycotoxin contamination when dairy animal consumed contaminated feed with mycotoxins. The most prevalent mycotoxins in milk are aflatoxin M₁ which is a good indicator of the level of aflatoxin B₁ in the dairy feed. Cheeses as milk product cannot escape mycotoxins contamination, especially aflatoxins by its affinity to bind to milk protein casein [9]. Furthermore, the direct growths of toxigenic fungi on a rich nutrient source as milk favour the production of mycotoxins and with in such commodities [10, 11].

Moreover, fungi are undesirable organisms which can grow at a wide range, causing spoilage, and are responsible for flavour defects, discolouration and reduced appearance of the product [12]. They were found as commensally in the gastrointestinal tract of human and animals. However, a wide variety of moulds have been involved in the infection of human, perhaps because of the intensive and prolonged use of antibiotics [13].

Up to date, nanotechnology is employed in various fields including food systems security, novel tools from molecular and cellular biology, new materials for pathogen detection and protection of the environment and food industry and safety [14]. Also, the advances in the field of nanotechnology have been used in industry, medicine and the pharmaceuticals and food packaging products [15]. Recently, chitosan nanoparticles (CSNPs) have been widely studied for the delivery of antibiotic drugs [16,17] and anti-cancer drugs, such as 5-fluorouracil, paclitaxel-doxorubicin, letrozole, and saponin [18, 19]. While, the antifungal potential of chitosan and chitosan nanoparticles (CSNPs) against *F. graminearum* colonies was detected and can be used as biopesticides [20], growth inhibitor for *Rhizopus* sp. and *A. niger* [21].

Therefore, the present study was undertaken

to recover the prevalence of fungi and aflatoxins in kareish cheese and evaluation the methods of detection of aflatoxins. Also, try to explore the role of chitosan nanoparticles in improvement the fungal quality and the physical properties of kareish cheese.

Material and Methods

Samples

A total of 40 collective kareish cheese samples about 250 g. of each were obtained from retail farmers and supermarkets from different district in Giza Governorate. The collected samples were transferred to the laboratory in clean, dry, sterile and tightly closed wide mouth Jars, with a minimum of delay, where they were prepared for the mycological examination and aflatoxins detection.

Skimmed raw milk: Fifteen kilogram of fresh buffalo's skimmed raw milk (0.5% fat) was obtained from a dairy farm in Giza.

Chitosan nanoparticles: It was obtained from the nanotechnology lab in Agriculture Research Centre (ARC), Giza, Egypt. The average particle size was 60-70 nm.

Isolation and Identification of Fungi in Cheese Samples

Enumeration and isolation of fungi according to ICMSF [22]:

Ten grams of each sample were transferred aseptically into a sterile blender jar, to which 90 ml of 1% peptone water were added and homogenized in a sterile warring blender for 2 minutes, and tenfold serial dilutions of the homogenate were prepared [22]. One millimeter quantities of the previously prepared serial dilutions were inoculated separately into Petri dish plates and mixed with Sabouraud dextrose agar medium. The plates were left to solidify after mixing and incubated at 25°C for 3-5 days. The counts of mould and yeast colonies were recorded. Individual suspected colonies were selected depending upon their morphological characters. The stock culture was made from each isolate and monitored on Czapek-Dox, malt extract and potato dextrose agar (PDA) slopes for further identification.

Identification of isolates

The colony count and identification of mould species was carried out by observation of macroscopic and microscopic characteristics of the fungi colonies according to ISO [23]

for genus *Aspergillus* and other genera were identified according to Pitt and Hocking [24] while the yeast isolates were identified according to the method of Van der Walt and Yarrow [25].

Detection of Aflatoxin M₁ in cheese samples

Estimation of aflatoxin M₁ by thin-layer chromatography (T.L.C.) It was proceeded according to the method described by AOAC [26].

Estimation of aflatoxin M₁ by a fluorometric immunoaffinity method

The positive samples for aflatoxin M₁ in the TLC method were measured quantitatively by the fluorometric method using specific FGisAfla test standards according to the recommended method of AOAC [27].

Experimental evaluation of the efficiency of chitosan nanoparticle in improving the characters of Kareish cheese.

Preparation of Kareish cheese: Fifteen kilograms of skimmed raw buffalo's milk were divided into 5 parts (each of 3 kg.) and every part subjected to preparation of cheese as recommended by Korish and Abdelhamid [28]. Kareish cheeses were manufactured as follows: skim milk (0.5% fat and 8.5% SNF) was heated at 74°C for 15 sec. and then rapidly cooled to 40°C. At this point yogurt starter culture was added at level of 1.5% to the base mixture for coagulation. After complete coagulation, the curd was separately transferred into gauze for wheing off in 24 hr., then cut and stored in its pasteurized salted whey (7% salt) for 24 hr. Resultant cheeses were stored at 4°C.

Cheese coating with chitosan nanoparticles [29]

Prepared kareish cheeses were divided into five parts each one was about 600 gram. In the first part, no additives were used and considered as control. While in the second and third parts, the used skim milk during manufacturing of cheese was immersed in nano-chitosan at the concentration of (0.25% and 0.5%). Whereas, in the fourth and fifth parts, after manufacturing of cheese, they were oated with chitosan nanoparticles by immersion in different concentration of (0.25% and 0.5%) up to 2 minutes. All parts were coated in sterile capped containers and stored at 4°C and 50 grams were examined every two days until the appearance of deterioration.

Mycological analyses of treated cheese [30]

The samples were analyzed for moulds and yeasts using sabouraud dextrose agar (SDA) on

the first day and every 48 hrs. Ten gram sample was taken aseptically from each cheese group and diluted 1:10 in sterile distilled water, and the mixture was homogenized for 2 min at 9000 rpm (Stomacher, Seward Medical, and the UK). After serial dilutions in PBS solution, aliquots of 100 µg were inoculated in selective media in duplicate by pour plate method. The plates were incubated at 25 ± 2°C for 2-5 days, to enumerate moulds and yeasts.

Evaluation of sensory properties of treated cheese with chitosan nanoparticles [31].

Kareish cheese samples of all batches about 40 gm were judged at zero-days and every 48 hours of refrigerated storage for taste, odour and colour which were evaluation. Taste, odour and colour were investigated on 10-point scale acceptability evaluation used a 10-point scale (1 = dislike very much, 5 = neither like nor dislike and 10 = like very much). Samples were presented in randomly coded glass plates.

Statistical analysis

Data obtained were analyzed statistically for descriptive statistics (mean, maximum, minimum and standard error) using SPSS 14 (2006) [32].

Results and Discussion

Occurrence of fungi in kareish cheese

Milk products as different types of cheese are most essential foodstuffs consumed by human and considered a source of high-quality animal protein having all the essential amino acids. It is also a rich source of calcium, phosphorus and many other micronutrients. All dairy products as cheese are probably most often spoiled by mould growth and may constitute public health hazards. While, Milk and milk products can quickly become moulded by a large number of species of fungi and a group of unicellular fungal organisms (yeast) during ripening, after cutting and slicing and during storage in shops or at home [28].

Recently, the role of microbes and their toxins in food safety gained colossal attention and high research activity. Not able to the prevalence of mycotoxigenic fungi and aflatoxins in food due to its toxicity, mutagenicity, teratogenicity and carcinogenicity [33].

In the present study, all examined kareish cheese samples were contaminated with fungi; the highest mean count cfu/g was (1X10³±0.4). The most common isolated mould species in fresh kareish cheese samples were *Aspergillus*

spp, which isolated from (80%). Regarding the incidence of members of *Aspergillus* species in milk and its products, *A.ochraceus*, *A. candidus* and *A.flavus* were recovered (45%, 45% and 20%) of kareish cheese samples. All samples were incriminated different yeast species. Several studies revealed that recovered different species of fungi from raw milk which represented six genera of moulds. The most prevalent fungi in these samples were the genus *Aspergillus* (60) with the mean of the count of $(1.6 \times 10^2 \pm 0.1)$ [34].

All examined kareish cheese samples were contaminated with yeast. The most predominant species of yeast was *C.albicans* which recovered (50%), followed by *Trichosporoncutaneum* (25%), *Cr.neoformans* and *Geotrichumcandidum* (12.5% for each) and *Torulopsis* sp. isolated from (7.5%) of kareish cheese samples, respectively.

In a similar study, [35] recovered several moulds from samples of cheese made from ewe

milk Spain. They were identified as *Penicillium* spp. in all cheese samples besides to other species, *Aspergillus*, *Geotrichum*, *Pullularia* (*Aureobasidium*), *Mucor*, *Paecilomyces*, *Candida* and *Acremonium*. However, Diaz et al. [36] isolated 24 strains of *Penicillium* and *Aspergillus* species from 22 samples of Spanish cheese and found that one strain only of *A. flavus* was toxigenic and capable of production of aflatoxin in M_1 .

On the other hand, Hassan et al. [1] recovered members of genus *Aspergillus* spp. from fresh Kareish cheese, fresh Damietta cheese and Yoghurt samples at rates of (70%, 65% and 60%) respectively. The species of *A. flavus* was recovered from 30% of Yoghurt, 25% of fresh Damietta cheese and 15% of fresh Kareish cheese, followed by *A. niger* which was recovered at the rates of (25%, 15%, and 30%) respectively.

TABLE 1. Prevalence and statistical analytical results of fungi in kareish cheese.

Fungal isolates	Incidence of Fungi		Count (cfu/g)		
	No.+ve	%	Min.	Max.	Mean \pm SE
Total fungi	40	100	1X10 ²	5X10 ⁷	1X10 ³ \pm 0.4
<i>Aspergillus</i> sp.	32	80	1X10	2X10 ⁷	1X10 ⁵ \pm 0.9
<i>A.flavus</i>	8	20	2X10 ³	1.5X10 ⁴	5X10 ³ \pm 0.99
<i>A.ochraceus</i>	18	45	1.0X10 ²	3X10 ³	1.5X10 ³ \pm 0.6
<i>A.candidus</i>	18	45	2X10 ¹	2.2X10 ⁴	2.5X10 ² \pm 0.45
Yeast species	40	100	2.4X10 ²	1X10 ⁴	3.7X10 ³ \pm 0.5

TABLE 2. Identification and statistical analytical of Yeasts that Recovered from Kareish Cheese.

Types of Yeast	Prevalence of yeast isolates		
	No. of +ve sample (40)	%	Count (cfu/g) Mean \pm SE
Yeast species	40	100	3.7X10 ³ \pm 0.5
<i>Candida albicans</i>	20	50	1.0X10 ² \pm 0.5
<i>Trichosporoncutaneum</i>	10	25	1.5X10 ¹ \pm 0.2
<i>Cr. Neoformans</i>	5	12.5	3.0X10 ² \pm 0.5
<i>Geotrichumcandidum</i>	5	12.5	2.1X10 ² \pm 0.5
<i>Torulopsis</i>	3	7.5	1.1X10 ² \pm 0.1
Total	43		

TABLE 3. Comparison between immune-affinity column and thin layer chromatography methods in detection of aflatoxins in kareish cheese.

Method of aflatoxin M_1 detection	Incidence of aflatoxin M_1 in samples		Levels of detected aflatoxins M_1 (PPb)(ug/kg)		
	No. of +ve	%	Min.	Max.	Mean \pm SE
Immune-affinity column	16	42%	2.4	5.9	0.414.2 \pm
Thin-layer chromatography	16	42%	1.1	2.5	1.7 \pm 0.11

*Food and Drug Administration (FDA) guidelines on the acceptable levels of aflatoxins in milk which has an action level of 0.5 ppb of aflatoxins.

Prevalence of aflatoxins in kareish cheese

In general, aflatoxins are known to be mutagenic, teratogenic, carcinogenic and immune suppressive in animals and possibly in man. This situation requires efficient analytical methods which are sensitive, quantitative and reliable to minimize the danger of these toxins through detection of the lowest quantities of Aflatoxin M₁ in food.

In the present study, the samples of kareish cheese were examined for detection of AFM₁ by thin-layer chromatography and immunoaffinity column methods. The residues of AFM₁ in 42 % were detected by both methods, but the sensitivity for higher quantitative detection was observed in immunoaffinity column than thin layer chromatography where the detection level ranges was higher in fluorometric method (2.4 ppb-5.9 ppb) with the mean levels of (4.2 ppb±0.41). While, in case of thin-layer chromatographic method, the detected levels ranged from (1.1 ppb - 2.5 ppb), with mean levels of (1.7 ppb±0.11). Hence, the fluorometric immunoaffinity column method is more effectively used for the detection of AFs in food than thin layer chromatography.

So, we concluded that the fluorometric method is more efficient than thin layer chromatography in detection of mycotoxins as aflatoxins.

Aflatoxin M₁ contamination of milk is considered as indicator of the level of aflatoxins B₁ and B₂ in feed, consequently cheese as milk product are able to be contaminated by their metabolic products M₁ and M₂, these findings are supported by many investigators [1,11,37, 38]. Several research studies detected aflatoxins B₁ and M₁ [39, 40] and aflatoxin M₁ only in the examined cheese samples [41-44].

The high levels of AFM₁ in kareish cheese in the present work could be attributed to the high moisture content of cheese and caused the growth of moulds which may assist the formation of mycotoxins in cheese [45]. Also, kareish cheese showed the highest level of aflatoxins contamination which may be attributed to the bad handling and long duration of storage.

Herein, the detection of aflatoxins in food by immunoaffinity column method is more accurate, sensitive and short time consuming than other old chromatographic methods. Hence in our study, the majority of kareish cheese samples had the aflatoxins levels more than 0.5 ppb and detected in high rate (42%), and this product

was rejected and considered unfit for human consumption. Similar findings were detected by several authors [1,11,46] who identified the ability of the fluorometric method to detect the lower levels of mycotoxins which not detected by the chromatographic method. Some researchers [47,48] established regulatory working guidelines on the acceptable levels of aflatoxins in milk which has an action level of 0.5 ppb of aflatoxins.

Hassan et al. [1] detected AFM₁ residues in fresh Kareish cheese over the permissible limits (0.5 ppb). Similarly, Ali, [11] detected AFs in milk and its products, the mean values of total aflatoxin residues (µg/kg) in fresh kareish cheese samples were (3.30 ± 2.21). The effects of aflatoxins in animal and human health varied according to the age, nutritional status, and amount of aflatoxins, frequency and composition of the diet [49]. The significant effects of Aflatoxins were mutagenic, carcinogenic, and teratogenic [50].

Therefore, numerous methods had been proposed to remove or inactivate microbial contamination in different types of food.

Influence of addition of Chitosan Nanoparticles to Kareish cheese during or after their preparations on survival of moulds and yeast count.

Recently, the nanomaterials are widely used in diversified sectors, including food systems. The Nano-sized particles have higher potential than their conventional sources, and thus they reduce the quantity required [51, 52]. They have more significant growth-promoting, immunomodulatory, antibacterial effects than conventional counterparts. The chitosan nanoparticle (CSNPs) have the amine groups change under weak acid condition to the -NH₃⁺ anion, which can interact with the cell wall of bacteria and hinder the growth of the microorganism [53].

In the present study, the addition of CSNPs before or after manufacturing of kareish cheese caused prolonged safe preservation of cheese as the CSNPs have the antimicrobial potential against several bacteria and fungi. Where the addition and coating of cheese with CSNPs prevent the contamination and growth of yeasts for more extended period than the control groups which not coated with chitosan nanoparticles (Table 4, Fig.1). When the added concentrations of CSNPs to kareish cheese increased the prolonged inhibition of fungal contamination and

colony count was decreased in comparison with the control cheese. Currently, the more antifungal potential was observed in case of addition of CSNPs after manufacturing of kareish cheese than in case of addition during ripening and preparation of kareish cheese. This is maybe due

to the contamination may occur after addition of CSNPs during preparation, but the addition after preparation overcome most fungal contamination. In general, both methods addition of CSNPs are efficient in the preservation and safe storage of kareish cheese for longer of time (Table 4, Fig.1).

TABLE 4. Influence of total fungi count (log cfu/g) after addition of Chitosan Nanoparticles to Kareish cheese during or after their preparations.

Prevalence of fungi in cheese (log cfu/g) before & after treatments with gradual concentrations of chitosan nanoparticles					Time of treatment (day)
0.5%		0.25%		0.0%	Time after treatments (days)
3 rd gp***	2 nd gp**	3 rd gp***	2 nd gp**	1 st gp*	
1.85	2.10	2.00	2.25	2.50	0
1.70	2.00	2.50	2.20	3.10	2
1.50	2.00	2.30	2.10	3.20	4
1.35	1.90	2.00	2.00	3.60	6
2.10	1.95	2.50	3.35	3.90	8
2.00	2.45	2.60	3.70	4.10	10
2.50	2.90	3.00	4.30	4.50	12
2.70	3.10	3.30	4.40	4.70	14
2.95	3.34	3.60	4.60	4.90	16
3.14	3.54	3.8	4.79	5.15	18
3.20	3.67	4.00	4.90	5.38	20

1stgp *= Control group, non-treated with CSNPs– 2ndgp **Second group (cheese treated during its preparation)- 3rd gp ***Third group (cheese treated after preparation).




Fig. 1. Influence of total fungi count (log cfu/g) after addition of chitosan nanoparticles to Kareish cheese during & after their preparations.

*Control group, Non-treated with nano chitosan - **Second group (cheese treated during its preparation) - ***Third group (cheese treated after preparation)

In a similar study, the CSNP have been used to improve the anti-bacterial effects [54]. Other research, detected that bio-nano composite of fish gelatin and chitosan nanoparticle have a good candidate for food packaging applications to the extent the shelf life of foods and products [55]. In addition, another research reported that CSNP exhibit higher antibacterial activity than chitosan solution [56]. Furthermore, El-Diasty et al. [2] revealed that the moulds and yeasts counts in case of treatment of kareishcheese with 1% chitosan were lower than control untreated cheese in which the fungal count increased during storage and reached to the high level by increase of storage period.

Evaluation of sensory properties of treated cheese with chitosan nanoparticles.

Some species of moulds are used in manufacturing and ripening of certain varieties of cheese even though other moulds are responsible for certain flavour defects and surface discolouration of cheese. Furthermore, some of these fungi produce toxic metabolites (mycotoxins) which may penetrate the cheese and polishing its surface might not sufficiently remove them and affect the health of consumers [1, 57].

In the present study, the colours, taste and odour of kareish cheese coated or added with CSNPs were improved more than 20 days and more accepted by the consumer, and palatable good characters were observed (Tables, 5, 6 and 7). Where, it is showed that the acceptability score evaluation was higher in treated cheese with CSNPs, mainly if coating occurred after the preparation of cheese and as the concentration

of CSNPs increased up to 0.5 %, the prolonged preservation and acceptability increased more than 20 days. While in the control group of untreated kareish cheese, the sensory properties (colour, taste and odour) were changed and loss its acceptability by consumers. Nearly similar, [2] detected that the treated cheesewith chitosan (0.5% and 1%) showed an improvement of shelf-life extended up to the 18th day of storage while in the control group of cheese, the changes of taste and texture were observed on the 6th day while the variations in colour appear by the 9th day.

From the obtained results, it becomes apparent that the addition of CSNPs to the Kareish cheese improves their preservation from fungal contamination changed their sensory properties to prolong their shelf-life. Moreover, CSNP coating after manufacture is more efficient in the preservation of kareish cheese than its addition to the cheese during production.

In spite of the different benefits of the application of nanotechnology in food products, the main focus, especially regarding milk and products coating. On the other hand, lack of knowledge about the impact of the nano material on human health is a significant obstacle for nano technology application, especially in food. Hence, more development of specific methods and models for valuation of the potential toxicity in nano structure materials is needed.

Therefore, extensive researches are still needed to support the effectiveness and safety of nano technology, avoid any harm to the environment or human beings and animals' health.

TABLE 5. Sensory colour property of Kareish cheese before and after addition of chitosan nanoparticles.

Time after treatments (days)	Changes in the colour of kareish cheese before and after treatments with gradual concentrations of Nano-chitosan				
	0.0%	0.25%		0.5%	
Colour	1 st gp *	2 nd gp **	3 rd gp ***	2 nd gp **	3 rd gp ***
0	10	10	10	10	10
2	10	10	10	10	10
4	10	10	10	10	10
6	5	10	10	10	10
8	5	10	10	10	10
10	1	10	10	10	10
12	1	5	10	10	10
14	1	5	10	10	10
16	1	5	5	5	10
18	1	1	5	5	5
20	1	1	1	5	5

1stgp *Control group, Non-treated with nano chitosan - 2ndgp ** (skim raw milk treated during its preparation) - 3rdgp *** (cheese treated after preparation). The overall acceptability score evaluation was a 10-point scale (1 = dislike very much, 5 = neither like nor dislike and 10 = like very much).

TABLE 6. Sensory odour property of Kareish cheese before and after addition of chitosan nanoparticles.

Time after treatments (days)	Changes in the odour of kareish cheese before and after treatments with gradual concentrations of Nano-chitosan				
	0.0%		0.25%		0.5%
	1 st gp *	2 nd gp **	3 rd gp ***	2 nd gp **	3 rd gp ***
0	10	10	10	10	10
2	10	10	10	10	10
4	10	10	10	10	10
6	10	10	10	10	10
8	5	10	10	10	10
10	1	10	10	10	10
12	1	5	10	10	10
14	1	5	10	10	10
16	1	5	10	10	10
18	1	1	5	5	10
20	1	1	5	5	10

1stgp * Control, Non-treated with nano chitosan -2ndgp ** (skim raw milk treated during its preparation) - 3rdgp *** (cheese treated after preparation). The overall acceptability score evaluation was a 10-point scale (1 = dislike very much, 5 = neither like nor dislike and 10 = like very much).

TABLE 7. Sensory taste property of Kareish cheese before and after addition of chitosan nanoparticles.

Time after treatment(days)	Changes in the taste of kareish cheese before and after treatments with gradual concentrations of Nano-chitosan				
	0.0%		0.25%		0.5%
	1 st gp *	2 nd gp **	3 rd gp ***	2 nd gp **	3 rd gp ***
0	10	10	10	10	10
2	10	10	10	10	10
4	10	10	10	10	10
6	5	10	10	10	10
8	1	7	10	10	10
10	1	7	10	10	10
12	1	5	10	10	10
14	1	5	10	10	10
16	1	5	8	5	10
18	1	1	5	5	10
20	1	1	5	5	10

1stgp *Control, non-treated with nano chitosan -2ndgp ** (skim raw milk treated during its preparation) - 3rdgp *** (cheese treated after preparation). The overall acceptability score evaluation was a 10-point scale (1 = dislike very much, 5 = neither like nor dislike and 10 = like very much).

Conclusion and Recommendations

Fungal contaminated milk products are widely distributed in nature and may lead to food poisoning which renders them unfit for consumption causing high economic losses, or they may be incriminated in mycotoxicosis to the consumers.

Under favourable environmental conditions (temperature, relative humidity, water activity, suitable medium and enough time), moulds

can grow well and produce their mycotoxins particularly aflatoxins, which are considered to be one of the most potent threats to human beings.

Thus, special care should be taken with manufacturing and storage practices of milk products. This could be achieved by proper handling of milk product, good manufacturing and storage. Also, chitosan nanoparticle is an effective natural antifungal preservative against yeasts and moulds at very low concentrations.

Cheese treated with chitosan nanoparticles could extend the shelf-life of during refrigeration storage period which is desired by manufacturers and consumers.

Although special attention is required for the toxicology of nanomaterials before nano technology application in the food system, hence further studies are urgently needed to address toxicological risks of the application of nano technology in food technology to support the effectiveness and safety of nanotechnology, avoid any harm to the environment or human beings and animals' health.

Acknowledgment: The authors are gratefully acknowledged to Prof. Dr. Atef, A. Hassan, Prof. of Mycology, Department of Mycology and Mycotoxins, Animal Health Research Institute Dokki, (ARC), for his continuous assistance and advice in initiating the achievement of this work.

Funds statement: The authors not gained any fund for this research.

Conflict of interest: The authors declare that there are no conflicts of interest.

Ethical consideration: Not required.

References

- Hassan, A.A., El-Mokhtar, Nahed M., El-Shafei, Howayda M. and Nosier, Shireen M. The proteolytic and lipolytic activity of fungi isolated from milk products with references to aflatoxins contamination. *Egypt. J. of Appl. Sci.*, **27**(12), 302-320(2012).
- El-Diasty, E.M., Nesreen, Z. Eleiwa and Hoda, A.M. Aida Using of Chitosan as Antifungal Agent in Kareish Cheese. *New York Science Journal*, **5**(9), 5-10(2012).
- Refai, M.K. and Hassan, Atef, A. Monograph on Mycotoxigenic Fungi and Mycotoxins in Foods and Feeds with Synopsis on studies done by authors on mycotoxigenic fungi and mycotoxins in foods and feeds (2013). (https://www.academia.edu/7688721-Atef_Hassan)
- Munimbazi, L. and Bullerman, B. High-Performance Liquid Chromatographic Method for the Determination of Moniliformin in Corn. *Journal of AOAC International*, **81**(5), 999-1004(1998).
- Refai, M.K., Abo El-Yazid, Heidy and Hassan, A.A. Aspergillus and Aspergillosis in man, animals and birds, A guide for classification and identification of Aspergilli, diseases caused by them, diagnosis and treatment (2014). https://www.academia.edu/7688721-Atef_Hassan, https://www.researchgate.net/publication/Atef_Hassan
- Fink-Gremmels, J. and Malekinejad, H. Clinical effects and biochemical mechanisms associated with exposure to the mycoestrogen zearalenone. *Animal Feed Science and Technology*, **137**(3-4), 326-341(2007).
- IARC "International Agency for Research on Cancer". Monographs on the Evaluation of Carcinogenic Risks to Human. Some Traditional Herbal Medicine, Some Mycotoxins, Naphthalene and Styrene; IARC: Lyon, France (2002).
- FAO "Food and Agriculture Organization". Global food losses and food waste: extent, causes and prevention. Rome (2011). (www.fao.org/fileadmin/user_upload/ags/publications/GFL_web.pdf)
- Stoloff, L. and Wood, G. Aflatoxin M₁ in manufactured dairy products produced in the United States in 1979. *J. Dairy Sci.*, **64**, 2426-2430(1981).
- Frazier, W.C. and Westhoff, D.C. *Food Microbiology*, 3rded., Frazier W.C. and Westhoff, D.C. (Ed.), McGraw-Hill Book, New York (1988).
- Ali, S.F. Studies on aflatoxigenic Aspergilli in meat, milk and their products. PhD. Thesis in Veterinary Medical Science (Mycology), Zagazik Univ., Egypt (2014).
- Walker, H.A. Spoilage of food by yeasts. *Journal of Food Technology*, **3**, 57-61(1977).
- Carter, G.R., and Cole, J.R. Diagnostic procedures. Veterinary Bacteriology and Mycology. 5ed, New York: Academic Press, Inc., Boston, Sydney, Tokyo, Toronto (1990).
- Chen, Y., Evans, J. and Feldlaufer, M. Horizontal and vertical transmission of viruses in the honey bee, *Apis mellifera*. *Journal of Invertebrate Pathology*, **92**(3), 152-159(2006).
- Gajjar, P., Pettee, B., Britt, D.W., Huang, W., Johnson, W.P., and Anderson, A.J. Antimicrobial activities of commercial nanoparticles against an environmental soil microbe, *Pseudomonas putida* KT2440. *Journal of Biological Engineering*, **3**(1), 9-14(2009).

16. Cover, N.F., Lai-Yuen, S., Parsons, A.K. and Kumar, A. Synergetic effects of doxycycline-loaded chitosan nanoparticles for improving drug delivery and efficacy. *International Journal of Nanomedicine*, **7**, 2411-2419(2012).
17. Zaki, N.M. and Hafez, M.M. Enhanced antibacterial effect of ceftriaxone sodium- loaded chitosan nanoparticles against intracellular *Salmonella typhimurium*. *AAPS Pharm. Sci. Tech.*, **13**, 411-421 (2012).
18. Saboktakin, M. R., Tabatabaie, R. M., Maharramov, A. and Ramazanov, M. A. Synthesis and in vitro evaluation of carboxymethyl starch-chitosan nanoparticles as drug delivery system to the colon. *International Journal of Biological Macromolecules*, **48**(3), 381-385(2011).
19. Rejinold, N.S., Muthunarayanan, M., Divyarani, V.V., Sreerekha, P.R., Chennazhi, K.P., Nair, S.V. and Jayakumar, R. Curcumin-loaded biocompatible thermoresponsive polymeric nanoparticles for cancer drug delivery. *Journal of Colloid and Interface Science*, **360**(1), 39-51(2011).
20. Kheiri, A., MoosawiJorf, S.A., Malhipour, A., Saremi, H. and Nikkhah, M. Application of chitosan and chitosan nanoparticles for the control of Fusarium head blight of wheat *Fusariumgraminearum* in vitro and greenhouse. *Int. J. Biol. Macromol.*, **93**(PtA), 1261-1272(2016).
21. Chookhongkha, N., Sopondilok, T. and Photchanachai, S. Effect of chitosan and chitosan nanoparticles on fungal growth and chilli seed quality. Proc Int. Conf. Postharvest Pest Dis Manag Export Horti Crops-PPDM2012 973,231-237(2012).
22. ICMSF "International Commission on Microbiological Specifications for Foods". Microbiology in foods, their significance and methods of enumeration, 2nd ed. University of Toronto Press, Canada (1978).
23. ISO (21527-1/2008) International organization for standardization. Microbiology of food and animal feeding stuffs-Horizontal method for the enumeration of yeasts and moulds- Part 1: Colony count technique in products with water activity greater than 0,95. Switzerland, Geneva (2008).
24. Pitt, J.I. and Hocking, A.D. Fungi and food spoilage. Springer Science & Business Media press. Academic Press, New York, London, Toronto, Montreal, Tokyo (2009).
25. Van der Walt and Yarrow, D. Methods for the isolation, maintenance, classification and identification of yeasts. In: Kreger van Rij, N.J.W. (Ed.), *The Yeasts: A Taxonomic Study*. 3rd ed. Elsevier, 45-104(2009).
26. AOAC (Association Official Analytical Chemists). *Official Methods of Analysis*. 15th ed., Assoc. of official Analytical chemists, Washington, D.C.(2000).
27. AOAC (Association of Official Analytical Chemists). *Official Methods of Analysis*, 16th ed. chapter 10, International, Gaithersburg, Inc. Suite 400, 2200 Wilson Boulevard Arlington, Virginia 22201, USA (1995).
28. Korish, M. and Abd Elhamid, A. M. Improving the textural properties of Egyptian kariesh cheese by addition of hydrocolloids. *International Journal of Dairy Technology*, **65**(2), 237-242(2012).
29. Kouchak, M., Batool, S.N. and Maryam, N. Preparation and evaluation of composite coat containing chitosan nano particles on cheese to improve the stability of product. *International Journal of Comprehensive Leading Research In Science*, **1**(1), 41-55(2015).(ISSN: 2455-4693).
30. FDA "Food and drug administration". FDA Analytical manual yeast and mould and mycotoxins, chapter 18th ed., AOAC, Washington, DC (2001).
31. Kim, H.Y., Yu-Tae, J., In-Hue, B. and Hae-Soo, K. Physicochemical and Sensory Properties of Nanopowdered Chitosan-Added Maribo Cheese during Ripening. *Korean J. Food Sci. An.*, **34** (1), 57- 64(2014).
32. SPSS 14 (2006), Statistical Package for Social Science, SPSS for windows Release 14.0.0, 12 June, 2006. Standard Version, Copyright SPSS Inc., 1989-2006, All Rights Reserved, Copyright © SPSS Inc. (2006).
33. IARC "International Agency for Research on Cancer". Monographs on the Evaluation of Carcinogenic Risks to Humans. Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins. 56th ed. Lyon, France: IARC Press, (1993).
34. Hassan, A.A, Mansour-Mogda, K., Snousi-Samira, A.M. and Hassan-Randa, A. Mycological, biochemical and histopathol studies on acute fusariotoxicosis in sheep. *Life Science Journal*, **7** (3), 49-57(2010c).

35. Arizcun, C., Itulatin, M., Salmeron, J. and Torre, P. Study of cheeses with designation of Roncal and Idiazabal made in Navarra (Spain). *Acta Alimentaria*, **247**, 69-71 (1996).
36. Diaz, L.T.M., Blanco, R. C., Arias, G. M., Fernandez, G. M. and Leper, G. M. Mycotoxins in two Spanish cheese varieties. *Int. J. Food Microbiol.*, **30** (3), 391-395 (1996).
37. Abou-Zeid, A. M., Hassan, A. A. and Ragheb, R. R. Mycological studies on hard (Roume) and skim milk soft cheese (kariesh) with quantitative evaluation of the existing mycotoxins. *4th Sci. Cong. Proc. April, 3-6, 1996, Egypt. Vet. Med. J., Giza*, **44** (2), 113-121 (1996).
38. Zinedine, A., González-Osnaya, L., Soriano, J. M., Moltó, J.C., Idrissi, L., and Manes, J. Presence of aflatoxin M₁ in pasteurized milk from Morocco. *International Journal of Food Microbiology*, **114**(1), 25-29 (2007).
39. Van Egmond, H.P. Aflatoxins in milk (pp. 365-381). Academic Press, San Diego, CA. (1994).
40. El-Sawi, N.M., El-Maghraby, O.M., Mahran, H.S., Abo-Gharib, M. A. Abnormal contamination of cottage cheese in Egypt. *Journal of Applied Animal Research*, **6** (1), 81-90 (1994).
41. Williams, B.C. In mycotoxins, Scott, P.M. (Ed.), A Canadian perspective. Publication NRCC, CNRC, Ottawa, Ontario, p.49 (1985).
42. Dragacci, S. and Freymy, J. M. Application of immunoaffinity column clean up to aflatoxin M₁ Determination and survey in cheese. *J. Food Protect.*, **59** (9), 1011-1013 (1996).
43. Pietri, A., Bertuzzi, P. and Piva, G. Aflatoxin M₁ occurrence samples of grana padano cheese. *Food Ada. Contam.*, **14** (4) 341- 344 (1997).
44. Aly, S.A.M. Studies on mycotoxins in milk and some dairy products. PhD Thesis, Hygiene and Control of milk & its products. Faculty of Vet. Med., Cairo University, Egypt (1999).
45. Bullerman, L.B., Schroder, L.L. and Park, Y.K. Formation and control of mycotoxins in foods. *J. Food Prot.*, **47** (8), 637- 646 (1984).
46. Tania, A., Becker-Algeri, D.C., Kennedy, B., Camila, S., Deisy, A.D. and Eliana Badiale-Furlo. Mycotoxins in Bovine Milk and Dairy Products: A Review. *J. Food Sci.*, **81** (3), 544-552 (2016).
47. Commission Directive 2003/100/EC of 31 October 2003: amending Annex 1 to directive 2002/32/EC of the European Parliament and the Council on undesirable substances in animal feed and animal product (1/11/2003). *Official J. of the European Union*, **1** (285), 33-37 (2003).
48. FDA "Food and Drug Administration". 18th ed., AOAC, Washington, DC., (2005).
49. Olsen, J.H., Dragster, L. and Autrup, H. Cancer risk and occupational exposure to aflatoxins in Denmark. *Br. J. Cancer*, **58**, 392- 396 (1988).
50. Betina, V. Mycotoxins, Chemical, Biological and Environmental Aspects. Elsevier science Publishers, Amsterdam, Holland (1989).
51. Patil, S.S., Kore, K.B. and Kumar, P. Nanotechnology and its applications in veterinary and animal science. *Vet. World*, **2**, 475-477 (2012).
52. Sri Sindhura, K., Selvam, P.P., Prasad, T.N.V. and Hussain, O.M. Synthesis, characterization and evaluation of effect of phyto-genic zinc nanoparticles on soil exo-enzymes. *Appl. Nanosci.*, **4**, 819-827 (2014).
53. Zheng, L.Y. and Zhu, J.F. Study on antimicrobial activity of chitosan with different molecular weights. *Carbohydrate Polymers*, **54**, 527-530 (2003).
54. Ding, Y.S., Shen, X.F., Sithambaram, S., Gomez, S., Kumar, R., Crisostomo, V. M. B., and Aindow, M. Synthesis and catalytic activity of cryptomelane-type manganese dioxide nanomaterials produced by a novel solvent-free method. *Chemistry of Materials*, **17**(21), 5382-5389 (2005).
55. Hosseini, S.F., Rezaei, M., Zandi, M. and Farahmandghavi, F. Fabrication of bio-nanocomposite films based on fish gelatin reinforced with chitosan nanoparticles. *Food Hydrocolloids*, **44**, 172-182 (2015).
56. Qi, L., Xu, Z., Jiang, X., Hu, C. and Zou, X. Preparation and antibacterial activity of chitosan nano particles. *Carbohydr. Res.*, **339**, 2693-2700 (2004).
57. Northolt, M. D., Van Egmond, H. P., Soentoro, P. and De Jll, E. Fungal growth and the presence of sterigmatocystin in hard cheese. *J. Assoc. of Anal. Chem.*, **63**, 115-119 (1980).

محاولة لتحسين جودة الجبن القريش باستخدام جسيمات الشيتوزان النانوية

رشا محمود حمزة سيد الأهل^١، نهي متولي الشناوي^٢ أو خالد ناجي^٣
^١قسم بحوث الفطريات و السموم الفطرية - معهد بحوث الصحة الحيوانية -مركز البحوث الزراعية - القاهرة - مصر.
^٢المعمل المرجعي لسلامة الأغذية- معهد بحوث الصحة الحيوانية - مركز البحوث الزراعية - القاهرة - مصر.
^٣ قسم هندسة و تعبئة الأغذية - معهد تكنولوجيا الغذاء -مركز البحوث الزراعية - القاهرة - مصر.

قد أجريت هذه الدراسة لتقييم إضافة نانو الشيتوزان إلى الجبن القريش لتحسين مواصفاتها ، من ناحية الرائحة والطعم واللون والجودة الميكروبية. تم تجميع ٤٠ عينة من الجبن القريش من البائعين في الأسواق ومحلات السوبر ماركت من احياء مختلفة في محافظة الجيزة ، كانت جميع العينات ملوثة بالفطريات. كان أعلى متوسط للعد (١ × ٢١٠ ± ٢,٤) لكل جرام و أكثر الأنواع شيوعا هو فطر الأسبرجيلس تم عزله بنسبه ٨٠٪ و جميع العينات كانت ملوثة بأنواع مختلفة من الخمائر. و قد كان ٤٢٪ من العينات بها سموم الافلاتوكسين م ١ وتم الكشف عن بقايا السموم بطريقه العمود المناعية التي كانت أفضل من طريقة الكروماتوجرافي. و لقد اتضح لنا ان اضافة او تغطية الجبن القريش قبل او بعد التصنيع بالنانو شيتوزان بنسبه (٢٥,٠٪ و ٥,٠٪) تسببت في الحفظ الآمن لفترة طويلة حيث ان النانو شيتوزان له القدرة المضادة للميكروبات ضد الكثير من الفطريات. و أيضا الخصائص الطبيعية للجبن المغلفة او المضاف اليها نانوشيتوزان من حيث اللون و الرائحة و الطعم كانت مقبولة و كذلك ادت الي إطالة مدة صلاحية الجبن القريش دون التأثير علي خواصها الفيزيائية و بالتالي فان هذه الدراسة توضح إمكانية تغليف الجبن القريش بالنانوشيتوزان حيث انها طبيعيه و قويه وآمنه و مضاد للفطريات لتحسين الجودة و زيادة مدة الصلاحية.